

Table of Contents

PRESCRIBING INFORMATION	3
HIGHLIGHTS OF PRESCRIBING INFORMATION	3
FULL PRESCRIBING INFORMATION: CONTENTS	3
FULL PRESCRIBING INFORMATION	4
1 INDICATIONS AND USAGE	4
2 DOSAGE AND ADMINISTRATION	4
2.1 Dosage Information.	4
2.2 Administration Instructions for Brivaracetam Oral Solution.	4
2.4 Discontinuation of Brivaracetam Oral Solution.	5
2.5 Patients with Hepatic Impairment	5
2.6 Co-administration with Rifampin	5
3 DOSAGE FORMS AND STRENGTHS.	5
4 CONTRAINDICATIONS	5
5 WARNINGS AND PRECAUTIONS	5
5.1 Suicidal Behavior and Ideation	5
5.2 Neurological Adverse Reactions.	6
5.3 Psychiatric Adverse Reactions.	7
5.4 Hypersensitivity: Bronchospasm and Angioedema	7
5.5 Serious Dermatologic Reactions.	7
5.6 Withdrawal of Antiepileptic Drugs	8
6 ADVERSE REACTIONS	8
6.1 Clinical Trials Experience.	8
6.2 Postmarketing Experience.	9
7 DRUG INTERACTIONS	10
7.1 Rifampin.	10
7.2 Carbamazepine.	10
7.3 Phenytoin	10
7.4 Levetiracetam	10
8 USE IN SPECIFIC POPULATIONS	10
8.1 Pregnancy	10
8.2 Lactation.	11
8.4 Pediatric Use.	11
8.5 Geriatric Use.	12
8.6 Renal Impairment	12

8.7 Hepatic Impairment.	12
9 DRUG ABUSE AND DEPENDENCE	12
9.1 Controlled Substance	12
9.2 Abuse	12
9.3 Dependence	12
10 OVERDOSAGE	12
11 DESCRIPTION.	13
12 CLINICAL PHARMACOLOGY	13
12.1 Mechanism of Action	13
12.2 Pharmacodynamics	13
12.3 Pharmacokinetics	14
13 NONCLINICAL TOXICOLOGY	17
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility	17
14 CLINICAL STUDIES.	18
16 HOW SUPPLIED/STORAGE AND HANDLING	20
16.1 How Supplied	20
16.2 Storage and Handling	20
17 PATIENT COUNSELING INFORMATION.	20
MEDICATION GUIDE	22

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use BRIVARACETAM ORAL SOLUTION safely and effectively. See full prescribing information for BRIVARACETAM ORAL SOLUTION.

BRIVARACETAM oral solution, CV

Initial U.S. Approval: 2016

RECENT MAJOR CHANGES

Warnings and Precautions (5.5)

8/2025

INDICATIONS AND USAGE

Brivaracetam oral solution is indicated for the treatment of partial-onset seizures in patients 1 month of age and older. (1)

DOSAGE AND ADMINISTRATION

- *Adults (16 Years and Older)*: The recommended starting dosage for monotherapy or adjunctive therapy is 50 mg twice daily (100 mg per day). Based on individual patient tolerability and therapeutic response, the dosage may be adjusted down to 25 mg twice daily (50 mg per day) or up to 100 mg twice daily (200 mg per day). (2.1)
- *Pediatric Patients (1 Month to less than 16 Years)*: The recommended dosage is based on body weight and is administered orally twice daily (2.1)
- *Hepatic Impairment*: Dose adjustment is recommended for all stages of hepatic impairment. (2.5)

DOSAGE FORMS AND STRENGTHS

- Oral solution: 10 mg/mL (3)

CONTRAINDICATIONS

Hypersensitivity to brivaracetam or any of the inactive ingredients in brivaracetam oral solution. (4)

WARNINGS AND PRECAUTIONS

- *Suicidal Behavior and Ideation*: Monitor patients for suicidal behavior and ideation. (5.1)
- *Neurological Adverse Reactions*: Monitor for somnolence and fatigue, and advise patients not to drive or operate machinery until

- they have gained sufficient experience on brivaracetam. (5.2)
- *Psychiatric Adverse Reactions*: Behavioral reactions including psychotic symptoms, irritability, depression, aggressive behavior, and anxiety; monitor patients for symptoms. (5.3)
- *Hypersensitivity: Bronchospasm and Angioedema*: Advise patients to seek immediate medical care. Discontinue and do not restart brivaracetam oral solution if hypersensitivity occurs. (5.4)
- *Serious Dermatologic Reactions*: Discontinue brivaracetam oral solution unless an alternative etiology is established (5.5)
- *Withdrawal of Antiepileptic Drugs*: Brivaracetam should be gradually withdrawn. (5.6)

ADVERSE REACTIONS

Adults: Most common adverse reactions (at least 5% for brivaracetam and at least 2% more frequently than placebo) are somnolence/sedation, dizziness, fatigue, and nausea/vomiting. (6.1)

Pediatric Patients: Most common adverse reactions are similar to those seen in adult patients. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Lupin Pharmaceuticals, Inc. at 1-800-399-2561 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- *Rifampin*: Because of decreased concentrations, increasing brivaracetam oral solution dosage in patients on concomitant rifampin is recommended. (2.6, 7.1)
- *Carbamazepine*: Because of increased exposure to carbamazepine metabolite, if tolerability issues arise, consider reducing carbamazepine dosage in patients on concomitant brivaracetam oral solution. (7.2)
- *Phenytoin*: Because phenytoin concentrations can increase, phenytoin levels should be monitored in patients on concomitant brivaracetam. (7.3)
- *Levetiracetam*: Brivaracetam had no added therapeutic benefit when co-administered with levetiracetam. (7.4)

USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data, may cause fetal harm. (8.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 09/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE
2	DOSAGE AND ADMINISTRATION
2.1	Dosage Information
2.2	Administration Instructions for Brivaracetam Oral Solution
2.4	Discontinuation of Brivaracetam Oral Solution
2.5	Patients with Hepatic Impairment
2.6	Co-administration with Rifampin
3	DOSAGE FORMS AND STRENGTHS
4	CONTRAINDICATIONS
5	WARNINGS AND PRECAUTIONS
5.1	Suicidal Behavior and Ideation
5.2	Neurological Adverse Reactions
5.3	Psychiatric Adverse Reactions
5.4	Hypersensitivity: Bronchospasm and Angioedema
5.5	Serious Dermatologic Reactions
5.6	Withdrawal of Antiepileptic Drugs
6	ADVERSE REACTIONS
6.1	Clinical Trials Experience
6.2	Postmarketing Experience
7	DRUG INTERACTIONS
7.1	Rifampin
7.2	Carbamazepine
7.3	Phenytoin
7.4	Levetiracetam

8	USE IN SPECIFIC POPULATIONS
8.1	Pregnancy
8.2	Lactation
8.4	Pediatric Use
8.5	Geriatric Use
8.6	Renal Impairment
8.7	Hepatic Impairment
9	DRUG ABUSE AND DEPENDENCE
9.1	Controlled Substance
9.2	Abuse
9.3	Dependence
10	OVERDOSAGE
11	DESCRIPTION
12	CLINICAL PHARMACOLOGY
12.1	Mechanism of Action
12.2	Pharmacodynamics
12.3	Pharmacokinetics
13	NONCLINICAL TOXICOLOGY
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
14	CLINICAL STUDIES
16	HOW SUPPLIED/STORAGE AND HANDLING
16.1	How Supplied
16.2	Storage and Handling
17	PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Brivaracetam oral solution is indicated for the treatment of partial-onset seizures in patients 1 month of age and older.

2 DOSAGE AND ADMINISTRATION

2.1 Dosage Information

Monotherapy or Adjunctive Therapy

The recommended dosage for patients 1 month of age and older is included in Table 1. In pediatric patients weighing less than 50 kg, the recommended dosing regimen is dependent upon body weight. When initiating treatment, gradual dose escalation is not required. Dosage should be adjusted based on clinical response and tolerability.

Table 1: Recommended Dosage for Patients 1 Month of Age and Older

Age and Body Weight	Initial Dosage	Minimum and Maximum Maintenance Dosage
Adults (16 years and older)	50 mg twice daily (100 mg per day)	25 mg to 100 mg twice daily (50 mg to 200 mg per day)
Pediatric patients weighing 50 kg or more	25 mg to 50 mg twice daily (50 mg to 100 mg per day)	25 mg to 100 mg twice daily (50 mg to 200 mg per day)
Pediatric patients weighing 20 kg to less than 50 kg	0.5 mg/kg to 1 mg/kg twice daily (1 mg/kg to 2 mg/kg per day)	0.5 mg/kg to 2 mg/kg twice daily (1 mg/kg to 4 mg/kg per day)
Pediatric patients weighing 11 kg to less than 20 kg	0.5 mg/kg to 1.25 mg/kg twice daily (1 mg/kg to 2.5 mg/kg per day)	0.5 mg/kg to 2.5 mg/kg twice daily (1 mg/kg to 5 mg/kg per day)
Pediatric patients weighing less than 11 kg	0.75 mg/kg to 1.5 mg/kg twice daily (1.5 mg/kg to 3 mg/kg per day)	0.75 mg/kg to 3 mg/kg twice daily (1.5 mg/kg to 6 mg/kg per day)

2.2 Administration Instructions for Brivaracetam Oral Solution

Brivaracetam can be initiated with oral administration.

Brivaracetam oral solution may be taken with or without food.

Brivaracetam Oral Solution

A calibrated measuring device is recommended to measure and deliver the prescribed dose accurately. A household teaspoon or tablespoon is not an adequate measuring device.

When using brivaracetam oral solution, no dilution is necessary. Brivaracetam oral solution may also be administered using a nasogastric tube or gastrostomy tube.

Discard any unused brivaracetam oral solution remaining after 5 months of first opening the bottle.

2.4 Discontinuation of Brivaracetam Oral Solution

Avoid abrupt withdrawal from brivaracetam oral solution in order to minimize the risk of increased seizure frequency and status epilepticus [see *Warnings and Precautions (5.6) and Clinical Studies (14)*].

2.5 Patients with Hepatic Impairment

The recommended dosage for patients with hepatic impairment is included in Table 2 [see *Use in Specific Populations (8.7) and Clinical Pharmacology (12.3)*].

Table 2: Recommended Dosage for Patients with Hepatic Impairment

Age and Body Weight	Initial Dosage	Maximum Maintenance Dosage
Adults (16 years and older)	25 mg twice daily (50 mg per day)	75 mg twice daily (150 mg per day)
Pediatric patients weighing 50 kg or more		
Pediatric patients weighing 20 kg to less than 50 kg	0.5 mg/kg twice daily (1 mg/kg per day)	1.5 mg/kg twice daily (3 mg/kg per day)
Pediatric patients weighing 11 kg to less than 20 kg	0.5 mg/kg twice daily (1 mg/kg per day)	2 mg/kg twice daily (4 mg/kg per day)
Pediatric patients weighing less than 11 kg	0.75 mg/kg twice daily (1.5 mg/kg per day)	2.25 mg/kg twice daily (4.5 mg/kg per day)

2.6 Co-administration with Rifampin

Increase the brivaracetam dosage in patients on concomitant rifampin by up to 100% (i.e., double the dosage) [see *Drug Interactions (7.1) and Clinical Pharmacology (12.3)*].

3 DOSAGE FORMS AND STRENGTHS

Oral Solution

- 10 mg/mL: slightly viscous, clear, colorless to yellowish, raspberry-flavored liquid.

4 CONTRAINDICATIONS

Hypersensitivity to brivaracetam or any of the inactive ingredients in brivaracetam oral solution (bronchospasm and angioedema have occurred) [see *Warnings and Precautions (5.4)*].

5 WARNINGS AND PRECAUTIONS

5.1 Suicidal Behavior and Ideation

Antiepileptic drugs (AEDs), including brivaracetam, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.

Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% CI:1.2, 2.7) of suicidal thinking or behavior compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behavior or ideation among 27,863 AED-treated

patients was 0.43%, compared to 0.24% among 16,029 placebo-treated patients, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients treated. There were four suicides in drug-treated patients in the trials and none in placebo-treated patients, but the number is too small to allow any conclusion about drug effect on suicide.

The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behavior beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5 to 100 years) in the clinical trials analyzed. Table 3 shows absolute and relative risk by indication for all evaluated AEDs.

Table 3: Risk of Suicidal Thoughts or Behaviors by Indication for Antiepileptic Drugs in the Pooled Analysis

Indication	Placebo Patients with Events Per 1000 Patients	Drug Patients with Events Per 1000 Patients	Relative Risk: Incidence of Events in Drug Patients/Incidence in Placebo Patients	Risk Difference: Additional Drug Patients with Events Per 1000 Patients
Epilepsy	1	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9

The relative risk for suicidal thoughts or behavior was higher in clinical trials in patients with epilepsy than in clinical trials in patients with psychiatric or other conditions, but the absolute risk differences were similar for the epilepsy and psychiatric indications.

Anyone considering prescribing brivaracetam oral solution or any other AED must balance the risk of suicidal thoughts or behaviors with the risk of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior. Should suicidal thoughts and behavior emerge during treatment, consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

5.2 Neurological Adverse Reactions

Brivaracetam causes somnolence, fatigue, dizziness, and disturbance in coordination. Patients should be monitored for these signs and symptoms and advised not to drive or operate machinery until they have gained sufficient experience on brivaracetam to gauge whether it adversely affects their ability to drive or operate machinery.

Somnolence and Fatigue

Brivaracetam causes dose-dependent increases in somnolence and fatigue-related adverse reactions (fatigue, asthenia, malaise, hypersomnia, sedation, and lethargy) [see *Adverse Reactions (6.1)*]. In the Phase 3 controlled adjunctive epilepsy trials, these events were reported in 25% of patients randomized to receive brivaracetam at least 50 mg/day (20% at 50 mg/day, 26% at 100 mg/day, and 27% at 200 mg/day) compared to 14% of patients who received placebo. The risk is greatest early in treatment but can occur at any time.

Dizziness and Disturbance in Gait and Coordination

Brivaracetam causes adverse reactions related to dizziness and disturbance in gait and coordination (dizziness, vertigo, balance disorder, ataxia, nystagmus, gait disturbance, and abnormal coordination) [see *Adverse Reactions (6.1)*]. In the Phase 3 controlled adjunctive epilepsy trials, these events were reported in 16% of patients randomized to receive brivaracetam at least 50 mg/day compared to 10% of patients who received placebo. The risk is greatest early in treatment but can occur at any time.

5.3 Psychiatric Adverse Reactions

Brivaracetam causes psychiatric adverse reactions. In the Phase 3 controlled adjunctive epilepsy trials, psychiatric adverse reactions were reported in approximately 13% of patients who received brivaracetam (at least 50 mg/day) compared to 8% of patients who received placebo. Psychiatric events included both non-psychotic symptoms (irritability, anxiety, nervousness, aggression, belligerence, anger, agitation, restlessness, depression, depressed mood, tearfulness, apathy, altered mood, mood swings, affect lability, psychomotor hyperactivity, abnormal behavior, and adjustment disorder) and psychotic symptoms (psychotic disorder along with hallucination, paranoia, acute psychosis, and psychotic behavior). A total of 1.7% of adult patients treated with brivaracetam discontinued treatment because of psychiatric reactions compared to 1.3% of patients who received placebo.

Psychiatric adverse reactions were also observed in open-label pediatric trials and were generally similar to those observed in adults [see *Adverse Reactions (6.1)* and *Use in Specific Populations (8.4)*].

5.4 Hypersensitivity: Bronchospasm and Angioedema

Brivaracetam can cause hypersensitivity reactions. Bronchospasm and angioedema have been reported in patients taking brivaracetam oral solution. If a patient develops hypersensitivity reactions after treatment with brivaracetam oral solution, the drug should be discontinued. Brivaracetam oral solution is contraindicated in patients with a prior hypersensitivity reaction to brivaracetam or any of the inactive ingredients [see *Contraindications (4)*].

5.5 Serious Dermatologic Reactions

Serious dermatologic reactions, including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), have been reported in patients treated with brivaracetam. Time to onset of the serious dermatologic reaction ranged from 3 to 45 days after brivaracetam initiation in reported cases. Brivaracetam oral solution should be discontinued at the first sign of a rash, unless the rash is clearly not drug-related. If signs or symptoms suggest a serious dermatologic reaction, use of brivaracetam oral solution should not be resumed and alternative therapy should be considered.

5.6 Withdrawal of Antiepileptic Drugs

As with most antiepileptic drugs, brivaracetam should generally be withdrawn gradually because of the risk of increased seizure frequency and status epilepticus [see *Dosage and Administration (2.4) and Clinical Studies (14)*]. But if withdrawal is needed because of a serious adverse event, rapid discontinuation can be considered.

6 ADVERSE REACTIONS

The following serious adverse reactions are described elsewhere in labeling:

- Suicidal Behavior and Ideation [see *Warnings and Precautions (5.1)*]
- Neurological Adverse Reactions [see *Warnings and Precautions (5.2)*]
- Psychiatric Adverse Reactions [see *Warnings and Precautions (5.3)*]
- Hypersensitivity: Bronchospasm and Angioedema [see *Warnings and Precautions (5.4)*]
- Serious Dermatologic Reactions [see *Warnings and Precautions (5.5)*]
- Withdrawal of Antiepileptic Drugs [see *Warnings and Precautions (5.6)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In all controlled and uncontrolled trials performed in adult epilepsy patients, brivaracetam was administered as adjunctive therapy to 2437 patients. Of these patients, 1929 were treated for at least 6 months, 1500 for at least 12 months, 1056 for at least 24 months, and 758 for at least 36 months. A total of 1558 patients (1099 patients treated with brivaracetam and 459 patients treated with placebo) constituted the safety population in the pooled analysis of Phase 3 placebo-controlled studies in patients with partial-onset seizures (Studies 1, 2, and 3) [see *Clinical Studies (14)*]. The adverse reactions presented in Table 4 are based on this safety population; the median length of treatment in these studies was 12 weeks. Of the patients in those studies, approximately 51% were male, 74% were Caucasian, and the mean age was 38 years.

In the Phase 3 controlled epilepsy studies, adverse events occurred in 68% of patients treated with brivaracetam and 62% treated with placebo. The most common adverse reactions occurring at a frequency of at least 5% in patients treated with brivaracetam doses of at least 50 mg/day and greater than placebo were somnolence and sedation (16%), dizziness (12%), fatigue (9%), and nausea and vomiting symptoms (5%).

The discontinuation rates due to adverse events were 5%, 8%, and 7% for patients randomized to receive brivaracetam at the recommended doses of 50 mg, 100 mg, and 200 mg/day, respectively, compared to 4% in patients randomized to receive placebo.

Table 4 lists adverse reactions for brivaracetam that occurred at least 2% more frequently for brivaracetam doses of at least 50 mg/day than placebo.

Table 4: Adverse Reactions in Pooled Placebo-Controlled Adjunctive Therapy Studies in Adult Patients with Partial-Onset Seizures (Brivaracetam 50 mg/day, 100 mg/day, and 200 mg/day)

Adverse Reactions	Brivaracetam(N=803) %	Placebo (N=459) %
Gastrointestinal disorders		
Nausea/vomiting symptoms	5	3
Constipation	2	0
Nervous system disorders		
Somnolence and sedation	16	8
Dizziness	12	7
Fatigue	9	4
Cerebellar coordination and balance disturbances*	3	1
Psychiatric disorders		
Irritability	3	1

* Cerebellar coordination and balance disturbances includes ataxia, balance disorder, coordination abnormal, and nystagmus.

There was no apparent dose-dependent increase in adverse reactions listed in Table 4 with the exception of somnolence and sedation.

Pediatric Patients

Safety of brivaracetam was evaluated in two open-label, safety and pharmacokinetic trials in pediatric patients 2 months to less than 16 years of age. Across studies of pediatric patients with partial onset seizures, 186 patients received brivaracetam oral solution or tablet, of whom 123 received brivaracetam for at least 12 months. Adverse reactions reported in clinical studies of pediatric patients were generally similar to those seen in adult patients. Decreased appetite was also observed in these pediatric trials.

Hematologic Abnormalities

Brivaracetam can cause hematologic abnormalities. In the Phase 3 controlled adjunctive epilepsy studies, a total of 1.8% of brivaracetam-treated patients and 1.1% of placebo-treated patients had at least one clinically significant decreased white blood cell count ($<3 \times 10^9/L$), and 0.3% of brivaracetam-treated patients and 0% of placebo-treated patients had at least one clinically significant decreased neutrophil count ($<1 \times 10^9/L$).

Comparison by Sex

There were no significant differences by sex in the incidence of adverse reactions.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of brivaracetam. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Skin and Subcutaneous Tissue Disorders: Serious dermatologic reactions (e.g., Stevens-Johnson syndrome and toxic epidermal necrolysis) [see *Warnings and Precautions (5.5)*].

7 DRUG INTERACTIONS

7.1 Rifampin

Co-administration with rifampin decreases brivaracetam plasma concentrations likely because of CYP2C19 induction [see *Clinical Pharmacology (12.3)*]. Prescribers should increase the brivaracetam dose by up to 100% (i.e., double the dosage) in patients while receiving concomitant treatment with rifampin [see *Dosage and Administration (2.6)*].

7.2 Carbamazepine

Co-administration with carbamazepine may increase exposure to carbamazepine-epoxide, the active metabolite of carbamazepine. Though available data did not reveal any safety concerns, if tolerability issues arise when co-administered, carbamazepine dose reduction should be considered [see *Clinical Pharmacology (12.3)*].

7.3 Phenytoin

Because brivaracetam can increase plasma concentrations of phenytoin, phenytoin levels should be monitored in patients when concomitant brivaracetam is added to or discontinued from ongoing phenytoin therapy [see *Clinical Pharmacology (12.3)*].

7.4 Levetiracetam

Brivaracetam provided no added therapeutic benefit to levetiracetam when the two drugs were co-administered [see *Clinical Studies (14)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Available data from the North American Antiepileptic Drug (NAAED) pregnancy registry, a prospective cohort study, case reports, and a case series are insufficient to identify a risk of major birth defects, miscarriage or other maternal or fetal outcomes associated with brivaracetam use during pregnancy. In animal studies, brivaracetam produced evidence of developmental toxicity (increased embryofetal mortality and decreased fetal body weights in rabbits; decreased growth, delayed sexual maturation, and long-term neurobehavioral changes in rat offspring) at maternal plasma exposures greater than clinical exposures [see *Data*].

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

Oral administration of brivaracetam (0, 150, 300, or 600 mg/kg/day) to pregnant rats during the period of organogenesis did not produce any significant maternal or embryofetal toxicity. The highest dose tested was associated with maternal plasma exposures (AUC) approximately 30 times exposures in humans at the maximum recommended dose (MRD) of 200 mg/day.

Oral administration of brivaracetam (0, 30, 60, 120, or 240 mg/kg/day) to pregnant rabbits during

the period of organogenesis resulted in embryofetal mortality and decreased fetal body weights at the highest dose tested, which was also maternally toxic. The highest no-effect dose (120 mg/kg/day) was associated with maternal plasma exposures approximately 4 times human exposures at the MRD.

When brivaracetam (0, 150, 300, or 600 mg/kg/day) was orally administered to rats throughout pregnancy and lactation, decreased growth, delayed sexual maturation (female), and long-term neurobehavioral changes were observed in the offspring at the highest dose. The highest no-effect dose (300 mg/kg/day) was associated with maternal plasma exposures approximately 7 times human exposures at the MRD.

Brivaracetam was shown to readily cross the placenta in pregnant rats after a single oral (5 mg/kg) dose of ¹⁴C-brivaracetam. From 1 hour post dose, radioactivity levels in fetuses, amniotic fluid, and placenta were similar to those measured in maternal blood.

8.2 Lactation

Risk Summary

Data from published literature indicate that brivaracetam is present in human milk. There is insufficient information on the effects of brivaracetam on the breastfed infant or on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for brivaracetam and any potential adverse effects on the breastfed infant from brivaracetam or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness of brivaracetam have been established in pediatric patients 1 month to less than 16 years of age. Use of brivaracetam in these age groups is supported by evidence from adequate and well-controlled studies of brivaracetam in adults with partial-onset seizures, pharmacokinetic data from adult and pediatric patients, and safety data in pediatric patients 2 months to less than 16 years of age [*see Dosage and Administration (2.1), Warnings and Precautions (5.3), Adverse Reactions (6.1), Clinical Pharmacology (12.3), and Clinical Studies (14)*].

Safety and effectiveness in pediatric patients below the age of 1 month have not been established.

Juvenile Animal Toxicity Data

The potential adverse effects of brivaracetam on postnatal growth and development were investigated in juvenile rats and dogs. Oral administration (0, 150, 300, or 600 mg/kg/day) to rats during the neonatal and juvenile periods of development (approximately equivalent to neonatal through adolescent development in humans) resulted in increased mortality, decreased body weight gain, delayed male sexual maturation, and adverse neurobehavioral effects at the highest dose tested and decreased brain size and weight at all doses. Therefore, a no-effect dose was not established; the lowest dose tested in juvenile rats was associated with plasma exposures (AUC) approximately 2 times those in children and adolescents at the recommended maintenance dose.

In dogs, oral administration (0, 15, 30, or 100 mg/kg/day) throughout the neonatal and juvenile periods of development induced liver changes similar to those observed in adult animals at the highest dose but produced no adverse effects on growth, bone density or strength, neurological testing, or neuropathology evaluation. The overall no-effect dose (30 mg/kg/day) and the no-effect dose for adverse effects on developmental parameters (100 mg/kg/day) were associated with plasma exposures approximately equal to and 4 times, respectively, those in children and adolescents at the recommended maintenance dose.

8.5 Geriatric Use

There were insufficient numbers of patients 65 years of age and older in the double-blind, placebo-controlled epilepsy trials (n=38) to allow adequate assessment of the effectiveness of brivaracetam in this population. In general, dose selection for an elderly patient should be judicious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy [see *Clinical Pharmacology* (12.3)].

8.6 Renal Impairment

Dose adjustments are not required for patients with impaired renal function. There are no data in patients with end-stage renal disease undergoing dialysis, and use of brivaracetam is not recommended in this patient population [see *Clinical Pharmacology* (12.3)].

8.7 Hepatic Impairment

Because of increases in brivaracetam exposure, dosage adjustment is recommended for all stages of hepatic impairment [see *Dosage and Administration* (2.5) and *Clinical Pharmacology* (12.3)].

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Brivaracetam oral solution contains brivaracetam and is listed as a Schedule V controlled substance.

9.2 Abuse

In a human abuse potential study, single doses of brivaracetam at therapeutic and supratherapeutic doses were compared to alprazolam (C-IV) (1.5 mg and 3 mg). Brivaracetam at the recommended single dose (50 mg) caused fewer sedative and euphoric effects than alprazolam; however, brivaracetam at supratherapeutic single doses (200 mg and 1000 mg) was similar to alprazolam on other measures of abuse.

9.3 Dependence

There was no evidence of physical dependence potential or a withdrawal syndrome with brivaracetam in a pooled review of placebo-controlled adjunctive therapy studies [see *Warnings and Precautions* (5.6)].

10 OVERDOSAGE


There is limited clinical experience with brivaracetam overdose in humans. Somnolence and dizziness were reported in a patient taking a single dose of 1400 mg (14 times the highest recommended single dose) of brivaracetam. The following adverse reactions were reported with

brivaracetam overdose: vertigo, balance disorder, fatigue, nausea, diplopia, anxiety, and bradycardia. In general, the adverse reactions associated with brivaracetam overdose were consistent with the known adverse reactions.

There is no specific antidote for overdose with brivaracetam. In the event of overdose, standard medical practice for the management of any overdose should be used. An adequate airway, oxygenation, and ventilation should be ensured; monitoring of cardiac rate and rhythm and vital signs is recommended. A certified poison control center should be contacted for updated information on the management of overdose with brivaracetam. There are no data on the removal of brivaracetam using hemodialysis, but because less than 10% of brivaracetam is excreted in urine, hemodialysis is not expected to enhance brivaracetam clearance.

11 DESCRIPTION

The chemical name of brivaracetam is (2S)-2-[(4R)-2-oxo-4-propyltetrahydro-1H-pyrrol-1-yl]butanamide. Its molecular formula is $C_{11}H_{20}N_2O_2$ and its molecular weight is 212.29. The chemical structure is:



Brivaracetam is a white to off-white crystalline powder. It is freely soluble in water, in ethanol, in methanol, in glacial acetic acid, in acetonitrile and in acetone. It is soluble in toluene.

Oral Solution

Brivaracetam oral solution contains 10 mg of brivaracetam per mL. The inactive ingredients are sodium citrate (trisodium citrate dihydrate), anhydrous citric acid, methylparaben, sodium carboxymethylcellulose, sucralose, non-crystallizing sorbitol solution, anhydrous glycerol, raspberry flavor, and purified water. Raspberry flavor also contains maltodextrin, propylene glycol and modified waxy maize starch.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The precise mechanism by which brivaracetam exerts its anticonvulsant activity is not known. Brivaracetam displays a high and selective affinity for synaptic vesicle protein 2A (SV2A) in the brain, which may contribute to the anticonvulsant effect.

12.2 Pharmacodynamics

Interactions with Alcohol

In a pharmacokinetic and pharmacodynamic interaction study in healthy subjects, co-administration of brivaracetam (single dose 200 mg [2 times greater than the highest recommended single dose]) and ethanol (continuous intravenous infusion to achieve a blood

alcohol concentration of 60 mg/100 mL during 5 hours) increased the effects of alcohol on psychomotor function, attention, and memory. Co-administration of brivaracetam and ethanol caused a larger decrease from baseline in saccadic peak velocity, smooth pursuit, adaptive tracking performance, and Visual Analog Scale (VAS) alertness, and a larger increase from baseline in body sway and in saccadic reaction time compared with brivaracetam alone or ethanol alone. The immediate word recall scores were generally lower for brivaracetam when co-administered with ethanol.

Cardiac Electrophysiology

At a dose 4 times the maximum recommended dose, brivaracetam did not prolong the QT interval to a clinically relevant extent.

12.3 Pharmacokinetics

Brivaracetam tablets, oral solution, and injection can be used interchangeably. Brivaracetam exhibits linear and time-independent pharmacokinetics at the approved doses.

The pharmacokinetics of brivaracetam are similar when used as monotherapy or as adjunctive therapy for the treatment of partial-onset seizures.

Absorption

Brivaracetam is highly permeable and is rapidly and almost completely absorbed after oral administration. Pharmacokinetics is dose-proportional from 10 to 600 mg (a range that extends beyond the minimum and maximum single-administration dose levels described in Dosage and Administration [*see Dosage and Administration (2.1)*]). The median T_{\max} for tablets taken without food is 1 hour (range 0.25 to 3 hours). Co-administration with a high-fat meal slowed absorption, but the extent of absorption remained unchanged. Specifically, when a 50 mg tablet was administered with a high-fat meal, C_{\max} (maximum brivaracetam plasma concentration during a dose interval, an exposure metric) was decreased by 37% and T_{\max} was delayed by 3 hours, but AUC (area under the brivaracetam plasma concentration versus time curve, an exposure metric) was essentially unchanged (decreased by 5%).

Distribution

Brivaracetam is weakly bound to plasma proteins ($\leq 20\%$). The volume of distribution is 0.5 L/kg, a value close to that of the total body water. Brivaracetam is rapidly and evenly distributed in most tissues.

Elimination

Metabolism

Brivaracetam is primarily metabolized by hydrolysis of the amide moiety to form the corresponding carboxylic acid metabolite, and secondarily by hydroxylation on the propyl side chain to form the hydroxy metabolite. The hydrolysis reaction is mediated by hepatic and extra-hepatic amidase. The hydroxylation pathway is mediated primarily by CYP2C19. In human subjects possessing genetic variations in CYP2C19, production of the hydroxy metabolite is decreased 2-fold or 10-fold, while the blood level of brivaracetam itself is increased by 22% or 42%, respectively, in individuals with one or both mutated alleles. CYP2C19 poor metabolizers and patients using inhibitors of CYP2C19 may require dose reduction. An additional hydroxy

acid metabolite is created by hydrolysis of the amide moiety on the hydroxy metabolite or hydroxylation of the propyl side chain on the carboxylic acid metabolite (mainly by CYP2C9). None of the 3 metabolites are pharmacologically active.

Excretion

Brivaracetam is eliminated primarily by metabolism and by excretion in the urine. More than 95% of the dose, including metabolites, is excreted in the urine within 72 hours after intake. Fecal excretion accounts for less than 1% of the dose. Less than 10% of the dose is excreted unchanged in the urine. Thirty-four percent of the dose is excreted as the carboxylic acid metabolite in urine. The terminal plasma half-life ($t_{1/2}$) is approximately 9 hours.

Specific Populations

Age

Pediatric Patients (2 months to less than 16 years): An open-label, single-arm, multicenter, pharmacokinetic study with a 3-week evaluation period and fixed 3-step up-titration using brivaracetam oral solution was conducted in 99 pediatric patients 2 months to less than 16 years of age. In those patients, plasma concentrations were shown to be dose-proportional. The pediatric pharmacokinetic profile for brivaracetam was determined in a population pharmacokinetic analysis using sparse plasma concentration data obtained in three open-label studies in 255 adult and pediatric patients with epilepsy 2 months to 22 years of age that received intravenous, oral solution, or oral tablet formulations.

A weight-based dosing regimen is necessary to achieve brivaracetam exposures in pediatric patients 1 month to less than 16 years of age that are similar to those observed in adults treated at effective doses of brivaracetam [see *Dosage and Administration (2.2)*]. The estimated plasma clearance was 1.09 L/h, 1.81 L/h, and 3.11 L/h for pediatric patients weighing 11 kg, 20 kg, and 50 kg, respectively. In comparison, plasma clearance was estimated at 3.58 L/h in adult patients (70 kg body weight).

Geriatric Population: In a study in elderly subjects (65 to 79 years old; creatinine clearance 53 to 98 mL/min/1.73 m²) receiving brivaracetam 200 mg twice daily (2 times the highest recommended dosage), the plasma half-life of brivaracetam was 7.9 hours and 9.3 hours in the 65 to 75 and >75 years groups, respectively. The steady-state plasma clearance of brivaracetam was slightly lower (0.76 mL/min/kg) than in young healthy controls (0.83 mL/min/kg).

Sex

There were no differences observed in the pharmacokinetics of brivaracetam between male and female subjects.

Race/Ethnicity

A population pharmacokinetic analysis comparing Caucasian and non-Caucasian patients showed no significant pharmacokinetic difference.

Renal Impairment

A study in adult subjects with severe renal impairment (creatinine clearance <30 mL/min/1.73m² and not requiring dialysis) revealed that the plasma AUC of brivaracetam was moderately increased (21%) relative to healthy controls, while the AUCs of the acid, hydroxy, and

hydroxyacid metabolites were increased 3-fold, 4-fold, and 21-fold, respectively. The renal clearance of these inactive metabolites was decreased 10-fold. Brivaracetam has not been studied in patients undergoing hemodialysis [see *Use in Specific Populations (8.6)*].

Hepatic Impairment

A pharmacokinetic study in adult subjects with hepatic cirrhosis, Child-Pugh grades A, B, and C, showed 50%, 57%, and 59% increases in brivaracetam exposure, respectively, compared to matched healthy controls. The effect of hepatic impairment on brivaracetam pharmacokinetics in pediatric patients is expected to be comparable to the effect observed in adults [see *Dosage and Administration (2.5)* and *Use in Specific Populations (8.7)*].

Drug Interaction Studies

In Vitro Assessment of Drug Interactions

Drug-Metabolizing Enzyme Inhibition

Brivaracetam did not inhibit CYP1A2, 2A6, 2B6, 2C8, 2C9, 2D6, or 3A4. Brivaracetam weakly inhibited CYP2C19 and would not be expected to cause significant inhibition of CYP2C19 in humans. Brivaracetam was an inhibitor of epoxide hydrolase, (IC₅₀ = 8.2 μM), suggesting that brivaracetam can inhibit the enzyme *in vivo*.

Drug-Metabolizing Enzyme Induction

Brivaracetam at concentrations up to 10 μM caused little or no change of mRNA expression of CYP1A2, 2B6, 2C9, 2C19, 3A4, and epoxide hydrolase. It is unlikely that brivaracetam will induce these enzymes *in vivo*.

Transporters

Brivaracetam was not a substrate of P-gp, MRP1, or MRP2. Brivaracetam did not inhibit or weakly inhibit BCRP, BSEP, MATE1, MATE2/K, MRP2, OAT1, OAT3, OCT1, OCT2, OATP1B1, OATP1B3, or P-gp, suggesting that brivaracetam is unlikely to inhibit these transporters *in vivo*.

In Vivo Assessment of Drug Interactions

Drug Interaction Studies with Antiepileptic Drugs (AEDs)

Potential interactions between brivaracetam (25 mg twice daily to 100 mg twice daily) and other AEDs were investigated in a pooled analysis of plasma drug concentrations from all Phase 2 and 3 studies and in a population exposure-response analysis of placebo-controlled, Phase 3 studies in adjunctive therapy in the treatment of partial-onset seizures. None of the interactions require changes in the dose of brivaracetam oral solution. Interactions with carbamazepine and phenytoin can be clinically important [see *Drug Interactions (7.2)* and *(7.3)*]. The interactions are summarized in Table 5.

Table 5: Drug Interactions Between Brivaracetam and Concomitant Antiepileptic Drugs

Concomitant AED	Influence of AED on Brivaracetam	Influence of Brivaracetam on AED
Carbamazepine	26% decrease in plasma concentration	None for carbamazepine Increase of carbamazepine-epoxide metabolite* [see <i>Drug Interactions (7.2)</i>]
Lacosamide	No data	None

Lamotrigine	None	None
Levetiracetam	None	None
Oxcarbazepine	None	None on the active monohydroxy metabolite derivative (MHD)
Phenobarbital	19% decrease in plasma concentration	None
Phenytoin	21% decrease in plasma concentration	Up to 20% increase in plasma concentration [see Drug Interactions (7.3)]**
Pregabalin	No data	None
Topiramate	None	None
Valproic acid	None	None
Zonisamide	No data	None

* Brivaracetam is a reversible inhibitor of epoxide hydrolase resulting in an increased concentration of carbamazepine epoxide, an active metabolite of carbamazepine. The carbamazepine epoxide plasma concentration increased up to 198% at a brivaracetam oral solution dose of 100 mg twice daily.

** At a supratherapeutic dose of 400 mg/day brivaracetam, there was a 20% increase in phenytoin plasma concentration.

Drug Interaction Studies with Other Drugs

Effect of Other Drugs on Brivaracetam

Co-administration with CYP inhibitors or transporter inhibitors is unlikely to significantly affect brivaracetam exposure.

Co-administration with rifampin decreases brivaracetam plasma concentrations by 45%, an effect that is probably the result of CYP2C19 induction [see Dosage and Administration (2.6) and Drug Interactions (7.1)].

Oral Contraceptives

Co-administration of brivaracetam 200 mg twice daily (twice the recommended maximum daily dosage) with an oral contraceptive containing ethinylestradiol (0.03 mg) and levonorgestrel (0.15 mg) reduced estrogen and progestin AUCs by 27% and 23%, respectively, without impact on suppression of ovulation. However, co-administration of brivaracetam 50 mg twice daily with an oral contraceptive containing ethinylestradiol (0.03 mg) and levonorgestrel (0.15 mg) did not significantly influence the pharmacokinetics of either substance. The interaction is not expected to be of clinical significance.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

In a carcinogenicity study in mice, oral administration of brivaracetam (0, 400, 550, or 700 mg/kg/day) for 104 weeks increased the incidence of liver tumors (hepatocellular adenoma and carcinoma) in male mice at the two highest doses tested. At the dose (400 mg/kg) not associated with an increase in liver tumors, plasma exposures (AUC) were approximately equal to those in humans at the maximum recommended dose (MRD) of 200 mg/day. Oral administration (0, 150, 230, 450, or 700 mg/kg/day) to rats for 104 weeks resulted in an increased incidence of thymus tumors (benign thymoma) in female rats at the highest dose tested. At the highest dose not associated with an increase in thymus tumors, plasma exposures were approximately 9 times those in humans at the MRD.

Mutagenesis

Brivaracetam was negative for genotoxicity in *in vitro* (Ames, mouse lymphoma, and CHO chromosomal aberration) and *in vivo* (rat bone marrow micronucleus) assays.

Impairment of Fertility

Oral administration of brivaracetam (0, 100, 200, or 400 mg/kg/day) to male and female rats prior to and throughout mating and early gestation produced no adverse effects on fertility. The highest dose tested was associated with plasma exposures approximately 6 (males) and 13 (females) times those in humans at the MRD.

14 CLINICAL STUDIES

The effectiveness of brivaracetam in partial-onset seizures with or without secondary generalization was established in 3 fixed-dose, randomized, double-blind, placebo-controlled, multicenter studies (Studies 1, 2, and 3), which included 1550 patients. Patients enrolled had partial-onset seizures that were not adequately controlled with 1 to 2 concomitant antiepileptic drugs (AEDs). In each of these studies, 72% to 86% of patients were taking 2 or more concomitant AEDs with or without vagal nerve stimulation. The median baseline seizure frequency across the 3 studies was 9 seizures per 28 days. Patients had a mean duration of epilepsy of approximately 23 years.

All trials had an 8-week baseline period, during which patients were required to have at least 8 partial-onset seizures. The baseline period was followed by a 12-week treatment period. There was no titration period in these studies. Study 1 compared doses of brivaracetam, 50 mg/day and 100 mg/day with placebo. Study 2 compared a dose of brivaracetam 50 mg/day with placebo. Study 3 compared doses of brivaracetam, 100 mg/day and 200 mg/day with placebo. Brivaracetam was administered in equally divided twice daily doses. Upon termination of brivaracetam oral solution treatment, patients were down-titrated over a 1-, 2-, and 4-week duration for patients receiving 25, 50, and 100 mg twice daily brivaracetam, respectively.

The primary efficacy outcome in Study 1 and Study 2 was the percent reduction in 7-day partial-onset seizure frequency over placebo, while the primary outcome for Study 3 was the percent reduction in 28-day partial-onset seizure frequency over placebo. The criteria for statistical significance for all 3 studies was $p < 0.05$. Table 6 presents the primary efficacy outcome of the percent change in seizure frequency over placebo, based upon each study's protocol-defined 7- and 28-day seizure frequency efficacy outcome.

Table 6: Percent Reduction in Partial-Onset Seizure Frequency over Placebo (Studies 1, 2, and 3)

	Percent Reduction Over Placebo (%)
STUDY 1^a	
Placebo (n=100)	-----
50 mg/day (n=99)	9.5
100 mg/day (n=100)	17
STUDY 2^a	

Placebo (n=96)	-----
50 mg/day (n=101)	16.9*
STUDY 3^b	
Placebo (n=259)	-----
100 mg/day (n=252)	25.2*
200 mg/day (n=249)	25.7*


* Statistically significant based on testing procedure with alpha = 0.05

^a Based upon 7-day seizure frequency

^b Based upon 28-day seizure frequency

Figure 1 presents the percentage of patients by category of reduction from baseline in partial-onset seizure frequency per 28 days for all pooled patients in the 3 pivotal studies. Patients in whom the seizure frequency increased are shown at left as “worse.” Patients with an improvement in percent reduction from baseline partial-onset seizure frequency are shown in the 4 right-most categories.

Figure 1: Proportion of Patients by Category of Seizure Response for Brivaracetam and Placebo Across all Three Double- Blind Trials



Treatment with Levetiracetam

In Studies 1 and 2, which evaluated brivaracetam dosages of 50 mg and 100 mg daily, approximately 20% of the patients were on concomitant levetiracetam. Although the numbers of patients were limited, brivaracetam provided no added benefit when it was added to levetiracetam.

Although patients on concomitant levetiracetam were excluded from Study 3, which evaluated 100 and 200 mg daily, approximately 54% of patients in this study had prior exposure to levetiracetam.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Brivaracetam Oral Solution

- 10 mg/mL is a slightly viscous, clear, colorless to yellowish, raspberry-flavored liquid. It is supplied in amber glass bottles:
300 mL bottles NDC 70748-411-01

16.2 Storage and Handling

Store at 25°C (77°F); excursions permitted between 15°C to 30°C (59°F to 86°F). See USP Controlled Room Temperature. Do not freeze brivaracetam oral solution.

Discard any unused brivaracetam oral solution remaining after 5 months of first opening the bottle.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide). The Medication Guide can be accessed on www.lupin.com/brivaracetamOralSol-mg.pdf, or by calling 1-800-399-2561.

Suicidal Behavior and Ideation

Counsel patients, their caregivers, and/or families that antiepileptic drugs, including brivaracetam oral solution, may increase the risk of suicidal thoughts and behavior, and advise patients to be alert for the emergence or worsening of symptoms of depression; unusual changes in mood or behavior; or suicidal thoughts, behavior, or thoughts about self-harm. Advise patients, their caregivers, and/or families to report behaviors of concern immediately to a healthcare provider [*see Warnings and Precautions (5.1)*].

Neurological Adverse Reactions

Counsel patients that brivaracetam causes somnolence, fatigue, dizziness, and gait disturbance. These adverse reactions, if observed, are more likely to occur early in treatment but can occur at any time. Advise patients not to drive or operate machinery until they have gained sufficient experience on brivaracetam to gauge whether it adversely affects their ability to drive or operate machinery [*see Warnings and Precautions (5.2)*].

Psychiatric Adverse Reactions

Advise patients that brivaracetam causes changes in behavior (e.g., aggression, agitation, anger, anxiety, and irritability) and psychotic symptoms. Instruct patients to report these symptoms immediately to their healthcare provider [*see Warnings and Precautions (5.3)*].

Hypersensitivity: Bronchospasm and Angioedema

Advise patients that symptoms of hypersensitivity including bronchospasm and angioedema can

occur with brivaracetam. Instruct them to seek immediate medical care should they experience signs and symptoms of hypersensitivity [see *Warnings and Precautions (5.4)*].

Serious Dermatologic Reactions

Advise patients of the early signs and symptoms of serious dermatologic adverse reactions and to report any occurrence immediately to a healthcare provider [see *Warnings and Precautions (5.5)*].

Withdrawal of Antiepileptic Drugs

Advise patients not to discontinue use of brivaracetam oral solution without consulting with their healthcare provider. Brivaracetam should normally be gradually withdrawn to reduce the potential for increased seizure frequency and status epilepticus [see *Warnings and Precautions (5.6)*].

Pregnancy

Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during brivaracetam therapy [see *Use in Specific Populations (8.1)*].

Lactation

Counsel patients that brivaracetam, the active ingredient in brivaracetam oral solution, is present in breast milk. Instruct patients to discuss with their healthcare provider if they are breastfeeding or intend to breastfeed [see *Use in Specific Populations (8.2)*].

Dosing Instructions

Counsel patients that brivaracetam oral solution may be taken with or without food [see *Dosage and Administration (2.2)*].

Advise patients that the dosage of brivaracetam oral solution should be measured using a calibrated measuring device and not a household teaspoon. Instruct patients to discard any unused brivaracetam oral solution after 5 months of first opening the bottle [see *Dosage and Administration (2.2)*].

LUPIN and the  are registered trademarks of Lupin Pharmaceuticals, Inc.

Manufactured for:

Lupin Pharmaceuticals, Inc.

Naples, FL 34108

United States.

Manufactured by:

Novel Laboratories, Inc.

400 Campus Drive

Somerset NJ 08873

United States.

Revised: September 2025

MEDICATION GUIDE
Brivaracetam (briv" a ra' se tam) CV
oral solution

What is the most important information I should know about brivaracetam oral solution?

Brivaracetam oral solution is a federally controlled substance (CV) because it can be abused or lead to dependence. Keep brivaracetam oral solution in a safe place to prevent misuse and abuse. Selling or giving away brivaracetam oral solution may harm others and is against the law.

Like other antiepileptic drugs, brivaracetam oral solution may cause suicidal thoughts or actions in a very small number of people, about 1 in 500 people taking it.

Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- new or worse depression
- feeling agitated or restless
- trouble sleeping (insomnia)
- acting aggressive, feeling angry, or being violent
- an extreme increase in activity and talking (mania)
- attempts to commit suicide
- new or worse anxiety
- panic attacks
- new or worse irritability
- acting on dangerous impulses
- other unusual changes in behavior or mood

Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

How can I watch for early symptoms of suicidal thoughts and actions?

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

Do not stop brivaracetam oral solution without first talking to a healthcare provider.

- Stopping brivaracetam oral solution suddenly can cause serious problems.
- Stopping a seizure medicine suddenly can cause seizures that will not stop (status epilepticus).

What is brivaracetam oral solution?

Brivaracetam oral solution is a prescription medicine used to treat partial-onset seizures in people 1 month of age and older. It is not known if brivaracetam oral solution is safe and effective in children younger than 1 month of age.

Who should not take brivaracetam oral solution?

Do not take brivaracetam oral solution if you are allergic to brivaracetam or any of the ingredients in brivaracetam oral solution. See the end of this Medication Guide for a complete list of ingredients in brivaracetam oral solution.

What should I tell my healthcare provider before starting brivaracetam oral solution?

Before taking brivaracetam oral solution, tell your healthcare provider about all of your medical conditions, including if you:

- have or had depression, mood problems, or suicidal thoughts or behavior.
- have liver problems.
- have abused or been dependent on prescription medicines, street drugs, or alcohol.
- are pregnant or plan to become pregnant. It is not known if brivaracetam oral solution will harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking brivaracetam oral solution. You and your healthcare provider will have to decide if you should take brivaracetam oral solution while you are pregnant.
- are breastfeeding or plan to breastfeed. Brivaracetam passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take brivaracetam oral solution.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Brivaracetam oral solution may affect the way other

medicines work, and other medicines may affect how brivaracetam oral solution works. Do not start a new medicine without first talking with your healthcare provider. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist each time you get a new medicine.

How should I take brivaracetam oral solution?

- Take brivaracetam oral solution exactly as your healthcare provider tells you.
- Your healthcare provider will tell you how much brivaracetam oral solution to take and when to take it.
- Your healthcare provider may change your dose if needed. Do not change your dose without talking to your healthcare provider.
- Take brivaracetam oral solution with or without food.
- If your healthcare provider has prescribed brivaracetam oral solution, be sure to ask your pharmacist for a medicine dropper or medicine cup to help you measure the correct amount of brivaracetam oral solution. Do not use a household teaspoon or tablespoon. Ask your pharmacist for instructions on how to use the measuring device the right way.
- If you take too much brivaracetam oral solution, call your Poison Control Center or go to the nearest emergency room right away.

What should I avoid while taking brivaracetam oral solution?

Do not drive or operate machinery until you know how brivaracetam oral solution affects you. Brivaracetam oral solution may cause drowsiness, tiredness, dizziness, and problems with your balance and coordination.

What are the possible side effects of brivaracetam oral solution?

Brivaracetam oral solution may cause serious side effects, including:

- See “**What is the most important information I should know about brivaracetam oral solution?**”
- **Nervous system problems.** Drowsiness, tiredness, and dizziness are common with brivaracetam oral solution, but can be severe. See “**What should I avoid while taking brivaracetam oral solution?**”. Brivaracetam oral solution can also cause problems with balance and coordination.
- **Mental (psychiatric) symptoms.** Brivaracetam oral solution can cause mood and behavior changes such as aggression, agitation, anger, anxiety, apathy, mood swings, depression, hostility, and irritability. Irritability and anxiety are common with brivaracetam oral solution, and can be severe. People who take brivaracetam oral solution can also get psychotic symptoms such as hallucinations (seeing or hearing things that are really not there), delusions (false or strange thoughts or beliefs), and unusual behavior.
- **Serious allergic reactions.** Stop taking brivaracetam oral solution and call your healthcare provider right away if you have any of these signs of a serious allergic reaction:
 - swelling of your face, mouth, lips, gums, tongue, throat, or neck
 - trouble breathing
- **Serious skin rashes.** Brivaracetam oral solution may cause a severe rash with blisters and peeling skin. This may occur around the mouth, nose, eyes, and genitals or over much of the body. A mild rash may become a serious reaction and may cause death. Call your healthcare provider right away if you have a rash, skin blisters, sores in the mouth, or hives.
Do not stop taking brivaracetam oral solution without first talking to your healthcare provider.

The most common side effects of brivaracetam oral solution in adults include:

- | | |
|--------------|-----------------------|
| • sleepiness | • feeling tired |
| • dizziness | • nausea and vomiting |

Side effects of brivaracetam oral solution in children 1 month to less than 16 years of age are similar to those seen in adults.

These are not all the possible side effects of brivaracetam oral solution. For more information, ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store brivaracetam oral solution?

- Store brivaracetam oral solution at room temperature between 59°F to 86°F (15°C to 30°C).
- Do not freeze brivaracetam oral solution.
- Safely throw away any opened bottle of brivaracetam oral solution after 5 months of first opening the bottle, even if there is medicine left in the bottle.

Keep brivaracetam oral solution and all medicines out of the reach of children.**General information about the safe and effective use of brivaracetam oral solution.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use brivaracetam oral solution for a condition for which it was not prescribed. Do not give brivaracetam oral solution to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about brivaracetam oral solution. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about brivaracetam oral solution that is written for health professionals.

What are the ingredients in brivaracetam oral solution?

Active ingredient: brivaracetam

Oral solution inactive ingredients: sodium citrate (trisodium citrate dihydrate), anhydrous citric acid, methylparaben, sodium carboxymethylcellulose, sucralose, non-crystallizing sorbitol solution, anhydrous glycerol, raspberry flavor, and purified water. Raspberry flavor also contains maltodextrin, propylene glycol and modified waxy maize starch.

Manufactured for:

Lupin Pharmaceuticals, Inc.

Naples, FL 34108

United States.

Manufactured by:

Novel Laboratories, Inc.

400 Campus Drive

Somerset NJ 08873

United States.



LUPIN and the  are registered trademarks of Lupin Pharmaceuticals, Inc.

For more information, go to www.lupin.com, or call 1-800-399-2561.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: September 2025