

Pentazocine and Naloxone Tablets, USP

Revised: August 2025



Analgesic for Oral Use Only

WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF PENTAZOCINE AND NALOXONE TABLETS

**Addiction, Abuse, and Misuse**  
Because the use of Pentazocine and Naloxone Tablets exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death, assess each patient's risk prior to prescribing and reassess all patients regularly for the development of these behaviors and conditions [see **Warnings**].

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of Pentazocine and Naloxone Tablets, especially during initiation or following a dosage increase. To reduce the risk of respiratory depression, proper dosing and titration of Pentazocine and Naloxone Tablets are essential [see **Warnings**].

Accidental Ingestion

Accidental ingestion of even one dose of Pentazocine and Naloxone Tablets, especially by children, can result in a fatal overdose of Pentazocine [see **Warnings**].

Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing of Pentazocine and Naloxone Tablets and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate [see **Warnings, Precautions, Drug Interactions**].

Neonatal Opioid Withdrawal Syndrome (NOMS)

Advise pregnant women using opioids for an extended period of time of the risk of Neonatal Opioid Withdrawal Syndrome, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery [see **Warnings**].

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)


Healthcare providers are strongly encouraged to complete a REMS compliant education program and to counsel patients and caregivers on serious risks, safe use, and the importance of reading the Medication Guide with each prescription [see **Warnings**].

DESCRIPTION


Pentazocine and Naloxone Tablets, USP contain pentazocine hydrochloride, USP, a partial opioid agonist, equivalent to 50 mg base and is a member of the benzazocine series (also known as the benzomorphans series), and naloxone hydrochloride, USP, an opioid antagonist equivalent to 0.5 mg base.

Pentazocine and Naloxone Tablets, USP are an analgesic for oral administration.

Chemically, pentazocine hydrochloride, USP is (2*R*\*, 6*R*\*, 11*R*\*)-1,2,3,4,5,6-Hexahydro-6,11-dimethyl-3-(3-methyl-2-butenyl)-2,6-methano-3-benzazocin-8-ol hydrochloride, a white, crystalline substance soluble in acidic aqueous solutions, and has the following structural formula:



Chemically, naloxone hydrochloride, USP is Morphinan-6-one, 4,5-epoxy-3,14-dihydroxy-17-(2-propenyl)-, hydrochloride, (5*α*,-). It is a slightly off-white powder, and is soluble in water and dilute acids, and has the following structural formula:



Inactive Ingredients: colloidal silicon dioxide, dibasic calcium phosphate, D&C Yellow No. 10, magnesium stearate, microcrystalline cellulose, sodium lauryl sulfate and pregelatinized starch.

CLINICAL PHARMACOLOGY

Mechanism of Action

Pentazocine is a mixed agonist-antagonist at opioid receptors. Pentazocine is a partial agonist at the mu opioid receptor and an agonist at the kappa opioid receptor.

Naloxone is an opioid antagonist.

Pharmacodynamics

Effects on the Central Nervous System

Pentazocine produces respiratory depression by direct action on brain stem respiratory centers. The respiratory depression involves a reduction in the responsiveness of the brain stem respiratory centers to both increases in carbon dioxide tension and electrical stimulation.

Pentazocine causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origins may produce similar findings). Marked mydriasis rather than miosis may be seen due to hypoxia in overdose situations.

Effects on the Gastrointestinal Tract and Other Smooth Muscle

Pentazocine causes a reduction in motility associated with an increase in smooth muscle tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone may be increased to the point of spasm, resulting in constipation. Other opioid-induced effects may include a reduction in biliary and pancreatic secretions, spasm of sphincter of Oddi, transient elevations in serum amylase, and opioid-induced esophageal dysfunction (OIED).

Effects on the Cardiovascular System

Pentazocine produces peripheral vasodilation which may result in orthostatic hypotension or syncope. Manifestations of histamine release and/or peripheral vasodilation may include pruritus, flushing, red eyes, sweating, and/or orthostatic hypotension.

Effects on the Endocrine System

Opioids inhibit the secretion of adrenocorticotropic hormone (ACTH), cortisol, and luteinizing hormone (LH) in humans [see **Adverse Reactions**]. They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon.

Use of opioids for an extended period of time may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The causal role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date [see **Adverse Reactions**].

Effects on the Immune System

Opioids have been shown to have a variety of effects on components of the immune system. The clinical significance of these findings is unknown. Overall, the effects of opioids appear to be modestly immunosuppressive.

Concentration-Efficacy Relationships

The minimum effective analgesic concentration will vary widely among patients, especially among patients who have been previously treated with opioid agonists. The minimum effective analgesic concentration of pentazocine for any individual patient may increase over time due to an increase in pain, the development of a new pain syndrome, and/or the development of analgesic tolerance [see **Dosage and Administration**].

Concentration-Adverse Reaction Relationships

There is a relationship between increasing pentazocine plasma concentration and increasing frequency of dose-related opioid adverse reactions such as nausea, vomiting, CNS effects, and respiratory depression. In opioid-tolerant patients, the situation may be altered by the development of tolerance to opioid-related adverse reactions [see **Dosage and Administration**].

Opioid Antagonist Effects

Pentazocine weakly antagonizes the analgesic effects of morphine, meperidine, and phenazocine; in addition, it produces incomplete reversal of cardiovascular, respiratory, and behavioral depression induced by morphine and meperidine. Pentazocine has about 1/50 the antagonistic activity of nalorphine. It also has sedative activity.

Naloxone when administered orally at 0.5 mg has no pharmacologic activity. Naloxone hydrochloride administered parenterally at the same dose is an antagonist to pentazocine and a pure antagonist to narcotic analgesics.

Pentazocine and Naloxone Tablets are a potent analgesic when administered orally. However, the presence of naloxone in Pentazocine and Naloxone Tablets is intended to prevent the effect of pentazocine if the product is misused by injection.

Studies in animals indicate that the presence of naloxone does not affect pentazocine analgesia when the combination is given orally. If the combination is given by injection the action of pentazocine is neutralized.

Pharmacokinetics

Onset of significant analgesia usually occurs between 15 and 30 minutes after oral administration, and duration of action is usually three hours or longer.

Pentazocine is well absorbed from the gastrointestinal tract. Concentrations in plasma coincide closely with the onset, duration, and intensity of analgesia. The time to mean peak concentration in 24 normal volunteers was 1.7 hours (range 0.5 to 4 hours) after oral administration and the mean plasma elimination half-life was 3.6 hours (range 1.5 to 10 hours).

Pentazocine is metabolized in the liver and excreted primarily in the urine. The products of the oxidation of the terminal methyl groups and glucuronide conjugates are excreted by the kidney. Elimination of approximately 80% of the total dose occurs within 24 hours. Pentazocine passes into the fetal circulation.

INDICATIONS AND USAGE

Pentazocine and Naloxone Tablets are indicated for the management of pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate.

Limitations of Use

Because of the risks of addiction, abuse, misuse, overdose, and death, which can occur at any dosage or duration, and persist over the course of therapy, reserve opioid analgesics, including Pentazocine and Naloxone Tablets, for use in patients for whom alternative treatment options are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.

CONTRAINDICATIONS

Pentazocine and Naloxone Tablets are contraindicated in patients with:

- Significant respiratory depression [see **WARNINGS**]
- Acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment [see **WARNINGS**]
- Patients with known or suspected gastrointestinal obstruction, including paralytic ileus [see **WARNINGS**]
- Patients with hypersensitivity to either pentazocine, naloxone, or any of the formulation excipients (e.g., anaphylaxis) [see **WARNINGS**].

WARNINGS

Addiction, Abuse, and Misuse

Pentazocine and Naloxone Tablets contain pentazocine, a Schedule IV controlled substance. As an opioid, Pentazocine and Naloxone Tablets expose users to the risks of addiction, abuse, and misuse [see **DRUG ABUSE AND DEPENDENCE**].

Although the risk of addiction in any individual is unknown, it can occur in patients appropriately prescribed Pentazocine and Naloxone Tablets. Addiction can occur at recommended dosages and if the drug is misused or abused. The risk of opioid-related overdose or overdose-related death is increased with higher opioid doses, and this risk persists over the course of therapy. In postmarketing studies, addiction, abuse, misuse, and fatal and non-fatal opioid overdose were observed in patients with long-term opioid use [ADVERSE REACTIONS; Postmarketing Experience].

Assess each patient's risk for opioid addiction, abuse, or misuse prior to prescribing Pentazocine and Naloxone Tablets, and reassess all patients receiving Pentazocine and Naloxone Tablets for the development of these behaviors and conditions. Risks are increased in patients with a personal or family history of substance abuse (including drug or alcohol abuse or addiction) or mental illness (e.g., major depression). The potential for these risks should not, however, prevent the proper management of pain in any given patient. Patients at increased risk may be prescribed opioids such as Pentazocine and Naloxone Tablets, but use in such patients necessitates intensive counseling about the risks and proper use of Pentazocine and Naloxone Tablets along with frequent reevaluation for signs of addiction, abuse, and misuse. Consider prescribing an opioid overdose reversal agent [see **WARNINGS, DOSAGE AND ADMINISTRATION**].

Opioids are sought for nonmedical use and are subject to diversion from legitimate prescribed use. Consider these risks when prescribing or dispensing Pentazocine and Naloxone Tablets. Strategies to reduce these risks include prescribing the drug in the smallest appropriate quantity and advising the patient on careful storage of the drug during the course of treatment and proper disposal of unused drug. Contact local state professional licensing board or state-controlled substances authority for information on how to prevent and detect abuse or diversion of this product.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression has been reported with the use of opioids, even when used as recommended. Respiratory depression, if not immediately recognized and treated, may lead to respiratory arrest and death. Management of respiratory depression may include close observation, supportive measures, and use of opioid antagonists, depending on the patient's clinical status [see **OVERDOSAGE**]. Carbon dioxide (CO<sub>2</sub>) retention from opioid-induced respiratory depression can exacerbate the sedating effects of opioids.

While serious, life-threatening, or fatal respiratory depression can occur at any time during the use of Pentazocine and Naloxone Tablets, the risk is greatest during the initiation of therapy or following a dosage increase.

To reduce the risk of respiratory depression, proper dosing and titration of Pentazocine and Naloxone Tablets are essential [see **DOSAGE AND ADMINISTRATION**]. Overestimating the Pentazocine and Naloxone Tablets dosage when converting patients from another opioid product can result in a fatal overdose with the first dose.

Accidental ingestion of even one dose of Pentazocine and Naloxone Tablets, especially by children, can result in respiratory depression and death due to an overdose of pentazocine.

Educate patients and caregivers on how to recognize respiratory depression and emphasize the importance of calling 911 or getting emergency medical help right away in the event of a known or suspected overdose [see **PRECAUTIONS, Information for Patients**].

**Patient Access to an Opioid Overdose Reversal Agent for the Emergency Treatment of Opioid Overdose**  
Inform patients and caregivers about opioid overdose reversal agents (e.g., naloxone, nalmeferi). Discuss the importance of having access to an opioid overdose reversal agent, especially if the patient has risk factors for overdose (e.g., concomitant use of CNS depressants, a history of opioid use disorder, or prior opioid overdose) or if there are household members (including children) or other close contacts at risk for accidental ingestion or opioid overdose. The presence of risk factors for overdose should not prevent the management of pain in any patient [see **WARNINGS**].

Discuss the options for obtaining an opioid overdose reversal agent (e.g., prescription, over-the-counter, or as part of a community-based program).

There are important differences among the opioid overdose reversal agents, such as route of administration, product strength, approved patient age range, and pharmacokinetics. Be familiar with these differences, as outlined in the approved labeling for those products, prior to recommending or prescribing such an agent.

Educate patients and caregivers on how to recognize respiratory depression, and how to use an opioid overdose reversal agent for the emergency treatment of opioid overdose. Emphasize the importance of calling 911 or getting emergency medical help, even if an opioid overdose reversal agent is administered [see **WARNINGS, OVERDOSAGE**].

Risks from Concomitant Use With Benzodiazepines or Other CNS Depressants

Profound sedation, respiratory depression, coma, and death may result from the concomitant use of Pentazocine and Naloxone Tablets with benzodiazepines and/or other CNS depressants, including alcohol (e.g., non-benzodiazepines, sedative/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antiepileptics, and other opioids). Because of these risks, reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate.

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioid analgesics alone. Because of similar pharmacological properties, it is reasonable to expect similar risk with the concomitant use of other CNS depressant drugs with opioid analgesics [see **PRECAUTIONS, Drug Interactions**].

If the decision is made to prescribe a benzodiazepine or other CNS depressant concomitantly with an opioid analgesic, prescribe the lowest effective dosages and minimum durations of concomitant use. In patients already receiving an opioid analgesic, prescribe a lower initial dose of the benzodiazepine or other CNS depressant than indicated in the absence of an opioid, and titrate based on clinical response. If an opioid analgesic is initiated in a patient already taking a benzodiazepine or other CNS depressant, prescribe a lower initial dose of the opioid analgesic, and titrate based on clinical response. Inform patients and caregivers of this potential interaction, educate them on the signs and symptoms of respiratory depression (including sedation).

If concomitant use is warranted, consider prescribing an opioid overdose reversal agent [see **WARNINGS, DOSAGE AND ADMINISTRATION, and OVERDOSAGE**].

Advise both patients and caregivers about the risks of respiratory depression and sedation when Pentazocine and Naloxone Tablets is used with benzodiazepines or other CNS depressants (including alcohol and illicit drugs). Advise patients not to drive or operate heavy machinery until the effects of concomitant use of the benzodiazepine or other CNS depressant have been determined. Screen patients for risk of substance use disorders, including opioid abuse and misuse, and warn them of the risk for overdose and death associated with the use of additional CNS depressants including alcohol and illicit drugs [see **PRECAUTIONS, Information for Patients, Drug Interactions**].

Neonatal Opioid Withdrawal Syndrome (NOMS)

Use of Pentazocine and Naloxone Tablets for an extended period of time during pregnancy can result in withdrawal in the neonate. Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. Observe newborns for signs of neonatal opioid withdrawal syndrome and manage accordingly. Advise pregnant women using opioids for an extended period of time of the risk of risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see **PRECAUTIONS, Information for Patients, Pregnancy**].

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a Risk Evaluation and Mitigation Strategy (REMS) for these products. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare providers are strongly encouraged to do all of the following:

- Complete a REMS-compliant education program offered by an accredited provider of continuing education (CE) or another education program that includes all the elements of the FDA Education Blueprint for Health Care Providers involved in the Management or Support of Patients with Pain.
- Discuss the safe use, serious risks, and proper storage and disposal of opioid analgesics with patients

and/or their caregivers every time these medicines are prescribed. The Patient Counseling Guide (PCG) can be obtained at this link: <http://www.fda.gov/OpioidAnalgesicREMSPCG>.

- Emphasize to patients and their caregivers the importance of reading the Medication Guide that they will receive from their pharmacist every time an opioid analgesic is dispensed to them.
- Consider using other tools to improve patient, household, and community safety, such as patient-prescriber agreements that reinforce patient-prescriber responsibilities.

To obtain further information on the opioid analgesic REMS and for a list of accredited REMS CME/CE, call 800-503-0784, or log on to [www.opioidanalgesicrems.com](http://www.opioidanalgesicrems.com). The FDA Blueprint can be found at [www.fda.gov/OpioidAnalgesicREMSBlueprint](http://www.fda.gov/OpioidAnalgesicREMSBlueprint).

Opioid-Induced Hyperalgesia and Allodynia

Opioid-Induced Hyperalgesia (OIH) occurs when an opioid analgesic paradoxically causes an increase in pain, or an increase in sensitivity to pain. This condition differs from tolerance, which is the need for increasing doses of opioids to maintain a defined effect [see **Dependence**]. Symptoms of OIH include (but may not be limited to) increased levels of pain upon opioid dosage increase, decreased levels of pain upon opioid dosage decrease, or pain from ordinarily non-painful stimuli (allodynia). These symptoms may suggest OIH only if there is no evidence of underlying disease progression, opioid tolerance, opioid withdrawal, or additive behavior.

Cases of OIH have been reported, both with short-term and longer-term use of opioid analgesics. Though the mechanism of OIH is not fully understood, multiple biochemical pathways have been implicated. Medical literature suggests a strong biological plausibility between opioid analgesics and OIH and allodynia. If a patient is suspected to be experiencing OIH, carefully consider appropriately decreasing the dose of the current opioid analgesic, or opioid rotation (safely switching the patient to a different opioid moiety) [see **Dosage and Administration, Warning**].

Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly, Cachectic, or Debilitated Patients

The use of Pentazocine and Naloxone Tablets in patients with acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment is contraindicated.

**Patients with Chronic Pulmonary Disease:** Pentazocine and naloxone-treated patients with significant chronic obstructive pulmonary disease or cor pulmonale, and those with a substantially decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory depression are at increased risk of decreased respiratory drive including apnea, even at recommended dosages of Pentazocine and Naloxone Tablets [see **WARNINGS**].

**Elderly, Cachectic, or Debilitated Patients:** Life-threatening respiratory depression is more likely to occur in elderly, cachectic, or debilitated patients because they may have altered pharmacokinetics or altered clearance compared to younger, healthier patients [see **WARNINGS**].

Regularly evaluate such patients, particularly when initiating and titrating Pentazocine and Naloxone Tablets and when Pentazocine and Naloxone Tablets are given concomitantly with other drugs that depress respiration [see **WARNINGS**]. Alternatively, consider the use of non-opioid analgesics in these patients.

Adrenal Insufficiency

Cases of adrenal insufficiency have been reported with opioid use, more often following greater than 1 month of use. Presentation of adrenal insufficiency may include non-specific symptoms and signs including nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. If adrenal insufficiency is suspected, confirm the diagnosis with diagnostic testing as soon as possible. If adrenal insufficiency is diagnosed, treat with physiologic replacement doses of corticosteroids. Wean the patient off of the opioid to allow adrenal function to recover and continue corticosteroid treatment until adrenal function recovers. Other opioids may be tried as some cases reported use of a different opioid without recurrence of adrenal insufficiency. The information available does not identify any particular opioids as being more likely to be associated with adrenal insufficiency.

Severe Hypotension

Pentazocine and Naloxone Tablets may cause severe hypotension including orthostatic hypotension and syncope in ambulatory patients. There is increased risk in patients whose ability to maintain blood pressure has already been compromised by a reduced blood volume or concurrent administration of certain CNS depressant drugs (e.g., phenothiazines or general anesthetics) [see **PRECAUTIONS, Information for Patients**]. Regularly evaluate these patients for signs of hypotension after initiating or titrating the dosage of Pentazocine and Naloxone Tablets. In patients with circulatory shock, Pentazocine and Naloxone Tablets may cause vasodilation that can further reduce cardiac output and blood pressure. Avoid the use of Pentazocine and Naloxone Tablets in patients with circulatory shock.

Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness

In patients who may be susceptible to the intracranial effects of CO<sub>2</sub> retention (e.g., those with evidence of increased intracranial pressure or brain tumors), Pentazocine and Naloxone Tablets may reduce respiratory drive, and the resultant CO<sub>2</sub> retention can further increase intracranial pressure. Evaluate such patients for signs of sedation and respiratory depression, particularly when initiating therapy with Pentazocine and Naloxone Tablets.

Opioids may also obscure the clinical course in a patient with a head injury. Avoid the use of Pentazocine and Naloxone Tablets in patients with impaired consciousness or coma.

Risk of Gastrointestinal Complications

Pentazocine and Naloxone Tablets are contraindicated in patients with known or suspected gastrointestinal obstruction, including paralytic ileus.

The administration of Pentazocine and Naloxone Tablets or other opioids may obscure the diagnosis or clinical course in patients with acute abdominal conditions.

Pentazocine and Naloxone Tablets may cause spasm of the sphincter of Oddi. Opioids may cause increases in serum amylase. Regularly evaluate patients with biliary tract disease, including acute pancreatitis, for worsening symptoms.

Cases of opioid-induced esophageal dysfunction (OIED) have been reported in patients taking opioids. The risk of OIED may increase as the dose and/or duration of opioids increases. Regularly evaluate patients for signs and symptoms of OIED (e.g., dysphagia, regurgitation, non-cardiac chest pain), and if necessary, adjust opioid therapy as clinically appropriate.

Increased Risk of Seizures in Patients with Seizure Disorders

The pentazocine in Pentazocine and Naloxone Tablets may increase the frequency of seizures in patients with seizure disorders, and may increase the risk of seizures occurring in other clinical settings associated with seizures. Regularly evaluate patients with a history of seizure disorders for worsened seizure control during Pentazocine and Naloxone Tablets therapy.

Withdrawal

Do not abruptly discontinue Pentazocine and Naloxone Tablets in a patient physically dependent on opioids. When discontinuing Pentazocine and Naloxone Tablets in a physically dependent patient, gradually taper the dosage. Rapid tapering of Pentazocine and Naloxone Tablets in a patient physically dependent on opioids may lead to a withdrawal syndrome and return of pain [see **Dosage and Administration, Drug Abuse and Dependence**]. Additionally, the use of Pentazocine and Naloxone Tablets, a mixed agonist/antagonist opioid analgesic, in patients who are receiving a full opioid agonist analgesic may reduce the analgesic effect and/or precipitate withdrawal symptoms. Avoid concomitant use of Pentazocine and Naloxone Tablets with a full opioid agonist analgesic.

Risks of Driving and Operating Machinery

Pentazocine and Naloxone Tablets may impair the mental or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery. Warn patients not to drive or operate dangerous machinery unless they are tolerant to the effects of Pentazocine and Naloxone Tablets and know how they will react to the medication [see **Patient Counseling Information**].

Acute CNS Manifestations

Patients receiving therapeutic doses of Pentazocine and Naloxone Tablets have experienced hallucinations (usually visual), disorientation, and confusion which have cleared spontaneously within a period of hours. The mechanism of this reaction is not known. Such patients should be very closely observed and vital signs checked. If the drug is reinstituted, it should be done with caution since these acute CNS manifestations may recur.

The amount of naloxone present in Pentazocine and Naloxone Tablets (0.5 mg per tablet) has no action when taken orally and will not interfere with the pharmacologic action of pentazocine. However, this amount of naloxone given by injection has profound antagonistic action to narcotic analgesics.

Severe, even lethal, consequences may result from misuse of tablets by injection either alone or in combination with other substances, such as pulmonary emboli, vascular occlusion, ulceration and abscesses, and withdrawal symptoms in narcotic dependent individuals..

PRECAUTIONS

Porphyria

Particular caution should be exercised in administering pentazocine to patients with porphyria since it may provoke an acute attack in susceptible individuals.

Cardiovascular Disease

Pentazocine can elevate blood pressure, possibly through the release of endogenous catecholamines. Particular caution should be exercised in conditions where alterations in vascular resistance and blood pressure might be particularly undesirable, such as in the acute phase of myocardial infarction. Pentazocine and Naloxone Tablets should be used with caution in patients with myocardial infarction who have nausea or vomiting.

Impaired Renal or Hepatic Function

Decreased metabolism of pentazocine by the liver in extensive liver disease may predispose to accentuation of side effects. Although laboratory tests have not indicated that pentazocine causes or increases renal or hepatic impairment, the drug should be administered with caution to patients with such impairment.

Biliary Surgery

Narcotic drug products are generally considered to elevate biliary tract pressure for varying periods following their administration. Some evidence suggests that pentazocine may differ from other marketed narcotics in

this respect (i.e., it causes little or no elevation in biliary tract pressures). The clinical significance of these findings, however, is not yet known.

INFORMATION FOR PATIENTS/CAREGIVERS

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Storage and Disposal

Because of the risks associated with accidental ingestion, misuse, and abuse, advise patients to store Pentazocine and Naloxone Tablets securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home. Inform patients that leaving Pentazocine and Naloxone Tablets unsecured can pose a deadly risk to others in the home [see **WARNINGS, DRUG ABUSE AND DEPENDENCE**].

Advise patients and caregivers that when medicines are no longer needed, they should be disposed of promptly. Expired, unwanted, or unused Pentazocine and Naloxone Tablets should be disposed of by flushing the unused medication down the toilet if a drug take-back option is not readily available. Inform patients that they can visit [www.fda.gov/drugdisposal](http://www.fda.gov/drugdisposal) for a complete list of medicines recommended for disposal by flushing, as well as additional information on disposal of unused medicines.

Addiction, Abuse, and Misuse

Inform patients that the use of Pentazocine and Naloxone Tablets, even when taken as recommended, can result in addiction, abuse, and misuse, which can lead to overdose and death [see **WARNINGS**]. Instruct patients not to share Pentazocine and Naloxone Tablets with others and to take steps to protect Pentazocine and Naloxone Tablets from theft or misuse.

Life-Threatening Respiratory Depression

Inform patients of the risk of life-threatening respiratory depression, including information that the risk is greatest when starting Pentazocine and Naloxone Tablets or when the dosage is increased, and that it can occur even at recommended dosages.

Educate patients and caregivers on how to recognize respiratory depression and emphasize the importance of calling 911 or getting emergency medical help right away in the event of a known or suspected overdose [see **WARNINGS, Life Threatening Respiratory Depression**].

Accidental Ingestion

Inform patients that accidental ingestion, especially by children, may result in respiratory depression or death [see **WARNINGS**].

Interactions with Benzodiazepines and Other CNS Depressants


Inform patients and caregivers that potentially fatal additive effects may occur if Pentazocine and Naloxone Tablets are used with benzodiazepines or other CNS depressants, including alcohol, and not to use these drugs concomitantly unless supervised by a healthcare provider [see **WARNINGS, PRECAUTIONS, Drug Interactions**].

Patient Access to an Opioid Overdose Reversal Agent for the Emergency Treatment of Opioid Overdose

Inform patients and caregivers about opioid overdose reversal agents (e.g., naloxone, nalmeferi). Discuss the importance of having access to an opioid overdose reversal agent, especially if the patient has risk factors for overdose (e.g., concomitant use of CNS depressants, a history of opioid use disorder, or prior opioid overdose) or if there are household members (including children) or other close contacts at risk for accidental ingestion or opioid overdose.

Discuss with the patient the options for obtaining an opioid overdose reversal agent (e.g., prescription, over-the-counter, or as part of a community-based program)

Pentazocine and Naloxone Tablets, USP



Concomitant use of monoamine oxidase inhibitors (MAOIs) with Pentazocine and Naloxone Tablets may cause CNS excitation and hypertension through their respective effects on catecholamines. Caution should therefore be observed in administering Pentazocine and Naloxone Tablets to patients who are currently receiving MAOIs or who have received them within the preceding 14 days.

Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics

Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics such as butorphanol, nalbuphine, pentazocine, buprenorphine, may reduce the analgesic effect of Pentazocine and Naloxone Tablets and/or precipitate withdrawal symptoms. Avoid concomitant use of these drugs.

Muscle Relaxants

The concomitant use of opioids and muscle relaxants may enhance the neuromuscular blocking action of skeletal muscle relaxants and produce an increased degree of respiratory depression.

Because respiratory depression may be greater than expected, decrease the dosage of Pentazocine and Naloxone Tablets and/or the muscle relaxant as necessary. Due to the risk of respiratory depression with concomitant use of skeletal muscle relaxants and opioids, consider prescribing an opioid overdose reversal agent [see **WARNINGS, DOSAGE AND ADMINISTRATION**].

Diuretics

Opioids can reduce the efficacy of diuretics by inducing the release of antidiuretic hormone.

Evaluate patients for signs of diminished diuresis and/or effects on blood pressure and increase the dosage of the diuretic as needed.

Anticholinergic Drugs

The concomitant use of anticholinergic drugs may increase risk of urinary retention and/or severe constipation, which may lead to paralytic ileus.

Evaluate patients for signs of urinary retention or reduced gastric motility when Pentazocine and Naloxone Tablets are used concomitantly with anticholinergic drugs.

Tobacco

Smoking tobacco could enhance the metabolic clearance rate of pentazocine reducing the clinical effectiveness of a standard dose of pentazocine.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term animal studies have not been completed to evaluate the carcinogenic potential of the combination or individual components of Pentazocine and Naloxone Tablets.

Mutagenesis

Studies to evaluate the mutagenic potential of the components of Pentazocine and Naloxone Tablets have not been conducted.

Impairment of Fertility

Studies in animals to evaluate the impact of Pentazocine and Naloxone Tablets on fertility have not been completed. The daily administration of 4 mg/kg to 20 mg/kg pentazocine subcutaneously to female rats during a 14 day pre-mating period and until the 13th day of pregnancy did not have any adverse effects on the fertility rate.

Infertility

Use of opioids for an extended period of time may cause reduced fertility in females and males of reproductive potential. It is not known whether these effects on fertility are reversible [see **ADVERSE REACTIONS**].

Pregnancy

Risk Summary

Use of opioid analgesics for an extended period of time during pregnancy may cause neonatal opioid withdrawal syndrome [see **Warnings and Precautions (5.3)**]. There are no available data with Pentazocine and Naloxone Tablets in pregnant women to inform a drug-associated risk for major birth defects and miscarriage. In animal reproduction studies, pentazocine administered subcutaneously to pregnant hamsters during the early gestational period produced neural tube defects (i.e., exencephaly and cranioschisis) at 2.6 times the maximum daily dose (MDD). In pregnant rats administered pentazocine:naloxone during organogenesis, there were increased incidences of resorptions and extra ribs at 0.2 times the MDD. There was no evidence of malformations in rats or rabbits [see Data]. Based on animal data, advise pregnant women of the potential risk to a fetus. The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Clinical Considerations

Fetal/Neonatal Adverse Reactions

Use of opioid analgesics for an extended period of time during pregnancy for medical or nonmedical purposes can result in physical dependence in the neonate and neonatal opioid withdrawal syndrome shortly after birth. Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high pitched cry, tremor, vomiting, diarrhea and failure to gain weight. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by the newborn. Observe newborns for symptoms of neonatal opioid withdrawal syndrome and manage accordingly [see **WARNINGS**].

Labor or Delivery

Opioids cross the placenta and may produce respiratory depression and psycho-physiologic effects in neonates. An opioid overdose reversal agent, such as naloxone or nalmefene, must be available for reversal of opioid-induced respiratory depression in the neonate. Pentazocine and Naloxone Tablets are not recommended for use in pregnant women during or immediately prior to labor, when other analgesic techniques are more appropriate. Opioid analgesics, including Pentazocine and Naloxone Tablets, can prolong labor through actions which temporarily reduce the strength, duration, and frequency of uterine contractions. However, this effect is not consistent and may be offset by an increased rate of cervical dilation, which tends to shorten labor. Monitor neonates exposed to opioid analgesics during labor for signs of excess sedation and respiratory depression.

Data

Animal Data

In a published report, a single dose of pentazocine administered to pregnant hamsters on Gestation Day 8 increased the incidence of neural tube defects (exencephaly and cranioschisis) at a dose of 196 mg/kg, SC (2.6-times the maximum daily human dose (MDD) of 600 mg/day pentazocine (12 tablets) on a mg/m<sup>2</sup> basis). No evidence of neural tube defects were reported following a dose of 98 mg/kg (1.3 times the MDD).

Animal reproduction studies testing the combination of pentazocine and naloxone during organogenesis have been completed in rats and rabbits. In rats, a pentazocine:naloxone dose of 64 mg/kg:0.64 mg/kg via oral gavage from Gestation Day 6 to 15 increased the incidences of resorptions and extra ribs (0.2 times the maximum daily human dose of pentazocine via 12 tablets on a mg/m<sup>2</sup> basis). There were no clear treatment related effects in rabbits treated from Gestation Day 6 to 18 with a pentazocine:naloxone dose of up to 64 mg/kg:0.64 mg/kg via oral gavage (0.3-times the maximum daily human dose of pentazocine via 12 tablets on a mg/m<sup>2</sup> basis).

Lactation

Risk Summary

Pentazocine is excreted in human milk. Caution should be exercised when Pentazocine and Naloxone Tablets are administered to a nursing woman.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Pentazocine and Naloxone Tablets and any potential adverse effects on the breastfed infant from Pentazocine and Naloxone Tablets or from the underlying maternal condition.

Clinical Considerations

Infants exposed to pentazocine and naloxone through breast milk should be monitored for excess sedation and respiratory depression. Withdrawal symptoms can occur in breastfed infants when maternal administration of an opioid analgesic is stopped, or when breast-feeding is stopped.

Pediatric Use

Safety and effectiveness in pediatric patients below the age of 12 years have not been established.

Geriatric Use

Elderly patients (aged 65 years or older) may have increased sensitivity to Pentazocine and Naloxone Tablets. In general, use caution when selecting a dosage for an elderly patient, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

Respiratory depression is the chief risk for elderly patients treated with opioids, and has occurred after large initial doses were administered to patients who were not opioid-tolerant or when opioids were co-administered with other agents that depress respiration. Titrate the dosage of Pentazocine and Naloxone Tablets slowly in geriatric patients and frequently reevaluate the patient for signs of central nervous system and respiratory depression [see **Warnings**].

Pentazocine and Naloxone are known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to regularly evaluate renal function.

ADVERSE REACTIONS

The following adverse reactions associated with the use of Pentazocine and Naloxone were identified in clinical studies or postmarketing reports. Because some of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

**Cardiovascular** - Hypertension, hypotension, circulatory depression, tachycardia, syncope.

**Respiratory** - Rarely, respiratory depression.

**Acute CNS Manifestations** - Hallucinations (usually visual), disorientation, and confusion.

**Other CNS Effects** - Grand mal convulsions, increase in intracranial pressure, dizziness, lightheadedness, hallucinations, sedation, euphoria, headache, confusion, disorientation; infrequently weakness, disturbed dreams, insomnia, syncope, and depression; and rarely tremor, irritability, excitement, tinnitus.

**Autonomic** - Sweating; infrequently flushing; and rarely chills.

**Gastrointestinal** - Nausea, vomiting, constipation, diarrhea, anorexia, dry mouth, biliary tract spasm, and

rarely abdominal distress.

**Allergic** - Edema of the face; anaphylactic shock; dermatitis, including pruritus; flushed skin, including plethora; infrequently rash, and rarely urticaria.

**Ophthalmic** - Visual blurring and focusing difficulty, miosis.

**Hematologic** - Depression of white blood cells (especially granulocytes), with rare cases of agranulocytosis, which is usually reversible, moderate transient eosinophilia.

**Dependence and Withdrawal Symptoms** - [See **WARNINGS, PRECAUTIONS, and DRUG ABUSE AND DEPENDENCE** Sections].

**Other** - Urinary retention, paresthesia, serious skin reactions, including erythema multiforme, Stevens-Johnson syndrome toxic epidermal necrolysis, and alterations in rate or strength of uterine contractions during labor.

- **Serotonin syndrome:** Cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of opioids with serotonergic drugs.
- **Adrenal insufficiency:** Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use.
- **Anaphylaxis:** Anaphylaxis has been reported with ingredients contained in Pentazocine and Naloxone Tablets.
- **Androgen deficiency:** Cases of androgen deficiency have occurred with use of opioids for an extended period of time. [see *Clinical Pharmacology*].
- **Hyperalgesia and Allodynia:** Cases of hyperalgesia and allodynia have been reported with opioid therapy of any duration [see *Warnings*].
- **Hypoglycemia:** Cases of hypoglycemia have been reported in patients taking opioids. Most reports were in patients with at least one predisposing risk factor (e.g., diabetes).
- **Opioid-induced esophageal dysfunction (OIED):** Cases of OIED have been reported in patients taking opioids, and may occur more frequently in patients taking higher doses of opioid, and/or in patients taking opioids longer term [see **WARNINGS**].

Adverse Reactions from Observational Studies

A prospective, observational cohort study estimated the risks of addiction, abuse, and misuse in patients initiating long-term use of Schedule II opioid analgesics between 2017 and 2021. Study participants included in one or more analyses had been enrolled in selected insurance plans or health systems for at least one year, were free of at least one outcome at baseline, completed a minimum number of follow-up assessments, and either: 1) filled multiple extended-release/long-acting opioid analgesic prescriptions during a 90 day period (n=978); or 2) filled any Schedule II opioid analgesic for at least 70 of 90 days (n=1,244). Those included also had no dispensing of the qualifying opioids in the previous 6 months.

Over 12 months:

- approximately 1% to 6% of participants across the two cohorts newly met criteria for addiction, as assessed with two validated interview-based measures of moderate-to-severe opioid use disorder based on Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) criteria, and
- approximately 9% and 22% of participants across the two cohorts newly met criteria for prescription opioid abuse and misuse [defined in *DRUG ABUSE AND DEPENDENCE*], respectively, as measured with a validated self-reported instrument.

A retrospective, observational cohort study estimated the risk of opioid involved overdose or opioid overdose-related death in patients with new long-term use of Schedule II opioid analgesics from 2006 through 2016 (n=220,249). Included patients had been enrolled in either one of two commercial insurance programs, one managed care program, or one Medicaid program for at least 9 months. New long-term use was defined as having Schedule II opioid analgesic prescriptions covering at least 70 days' supply over the 3 months prior to study entry and none during the preceding 6 months. Patients were excluded if they had an opioid-involved overdose in the 9 months prior to study entry. Overdose was measured using a validated medical code-based algorithm with linkage to the National Death Index database. The 5-year cumulative incidence estimates for opioid-involved overdose or opioid overdose-related death ranged from approximately 1.5% to 4% across study sites, counting only the first event during follow-up. Approximately 17% of opioid-involved overdoses observed over the entire study period (5-11 years, depending on the study site) were fatal. Higher baseline opioid dose was the strongest and most consistent predictor of opioid-involved overdose or opioid overdose-related death. Study exclusion criteria may have selected patients at lower risk of overdose, and substantial loss to follow-up (approximately 80%) also may have biased estimates.

The risk estimates from the studies described above may not be generalizable to all patients receiving opioid analgesics, such as those with exposures shorter or longer than the duration evaluated in the studies.

**To report SUSPECTED ADVERSE REACTIONS, contact Lupin Pharmaceuticals Inc. at 1-866-403-7592 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

DRUG ABUSE AND DEPENDENCE

Controlled Substance

Pentazocine and Naloxone Tablets contain pentazocine, a Schedule IV controlled substance.

Abuse

Pentazocine and Naloxone Tablets contains Pentazocine, a substance with a high potential for misuse and abuse, which can lead to the development of substance use disorder, including addiction [see **Warnings**].

Misuse is the intentional use, for therapeutic purposes, of a drug by an individual in a way other than prescribed by a healthcare provider or for whom it was not prescribed.

Abuse is the intentional, non-therapeutic use of a drug, even once, for its desirable psychological or physiological effects.

Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that may include a strong desire to take the drug, difficulties in controlling drug use (e.g., continuing drug use despite harmful consequences, giving a higher priority to drug use than other activities and obligations), and possible tolerance or physical dependence.

Misuse and abuse of Pentazocine and Naloxone Tablets increases a risk of overdose, which may lead to central nervous system and respiratory depression, hypotension, seizures, and death. The risk is increased with concurrent abuse of Pentazocine and Naloxone Tablets with alcohol and other CNS depressants. Abuse or addiction to opioids in some individuals may be accompanied by concurrent tolerance and symptoms of physical dependence of opioid. In addition, abuse of opioids can occur in the absence of addiction.

All patients treated with opioids require careful and frequent reevaluation for signs of misuse, abuse, and addiction, because use of opioid analgesic products carries the risk of addiction even under appropriate medical use. Patients at high risk of Pentazocine and Naloxone Tablets abuse include those with a history of prolonged use of any opioid, including products containing Pentazocine, those with a history of drug or alcohol abuse, or those who use Pentazocine and Naloxone Tablets in combination with other abused drugs.

"Drug-seeking" behavior is very common in persons with substance use disorders. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing, or referral, repeated "loss" of prescriptions, tampering with prescriptions, and reluctance to provide prior medical records or contact information for other treating healthcare provider(s). "Doctor shopping" (visiting multiple prescribers to obtain additional prescriptions) is common among people who abuse drugs and people with substance use disorder. Preoccupation with achieving adequate pain relief can be appropriate behavior in a patient with inadequate pain control.

Pentazocine and Naloxone Tablets, like other opioids, can be diverted for nonmedical use into illicit channels of distribution. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests, as required by state and federal law, is strongly advised.

Proper assessment of the patient, proper prescribing practices, periodic reevaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

Risks Specific to Abuse of Pentazocine and Naloxone Tablets

Abuse of Pentazocine and Naloxone Tablets poses a risk of overdose and death. The risk is increased with concurrent use of Pentazocine and Naloxone Tablets with alcohol and/or other CNS depressants.

Parental drug abuse is commonly associated with transmission of infectious diseases such as hepatitis and HIV.

Dependence

Both tolerance and physical dependence can develop during use of opioid therapy.

Tolerance is a physiological state characterized by a reduced response to a drug after repeated administration (i.e., a higher dose of a drug is required to produce the same effect that was once obtained at a lower dose).

Physical dependence is a state that develops as a result of a physiological adaptation in response to repeated drug use, manifested by withdrawal signs and symptoms after abrupt discontinuation or a significant dose reduction of a drug.

Withdrawal may be precipitated through the administration of drugs with opioid antagonist activity (e.g., naloxone, nalmefene) mixed agonist/antagonist analgesics (e.g., pentazocine, butorphanol, nalbuphine), and partial agonists (e.g., buprenorphine). Physical dependence may not occur to a clinically significant degree until after several days to weeks of continued use.

Do not abruptly discontinue Pentazocine and Naloxone Tablets in a patient physically dependent on opioids. Rapid tapering of Pentazocine and Naloxone Tablets in a patient physically dependent on opioids may lead to serious withdrawal symptoms, uncontrolled pain, and suicide. Rapid discontinuation has also been associated with attempts to find other sources of opioid analgesics, which may be confused with drug-seeking for abuse.

When discontinuing Pentazocine and Naloxone Tablets, gradually taper the dosage using a patient-specific plan that considers the following: the dose of Pentazocine and Naloxone Tablets the patient has been taking, the duration of treatment, and the physical and psychological attributes of the patient. To improve the likelihood of a successful taper and minimize withdrawal symptoms, it is important that the opioid tapering schedule is agreed upon by the patient. In patients taking opioids for an extended period of time at high doses, ensure that a multimodal approach to pain management, including mental health support (if needed), is in place prior to initiating an opioid analgesic taper [see *Dosage and Administration, and Warnings*].

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal signs [see *Pregnancy*].

OVERDOSAGE

Clinical Presentation

Acute overdose with Pentazocine and Naloxone Tablets can be manifested by respiratory depression, somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and, in some cases, pulmonary edema, bradycardia, hypotension, hypoglycemia, partial or complete airway obstruction, atypical snoring, and death. Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations [see **CLINICAL PHARMACOLOGY**]. Toxic leukoencephalopathy has been reported after opioid overdose and can present hours, days, or weeks after apparent recovery from the initial intoxication.

For pentazocine alone in single doses above 60 mg there have been reports of the occurrence of nalorphine-like psychotomimetic effects such as anxiety, nightmares, strange thoughts, and hallucinations. Somnolence, marked respiratory depression associated with hypertension and tachycardia have also been reported as have seizures, hypotension, hypoglycemia, dizziness, nausea, vomiting, lethargy, and paresthesias. The respiratory depression is antagonized by naloxone (see Treatment). Circulatory failure and deepening coma may occur in more severe cases, particularly in patients who have also ingested other CNS depressants such as alcohol, sedative/hypnotics, or antihistamines."

Treatment of Overdose

In case of overdose, priorities are the reestablishment of a patent and protected airway and institution of assisted or controlled ventilation, if needed. Employ other supportive measures (including oxygen and vasopressors) in the management of circulatory shock and pulmonary edema as indicated. Cardiac arrest or arrhythmias will require advanced life-support measures.

For clinically significant respiratory or circulatory depression secondary to opioid overdose, administer an opioid overdose reversal agent such as naloxone or nalmefene. As pentazocine is a mixed opioid agonist/antagonist, larger doses of naloxone or nalmefene may be needed to reverse the effects of an overdose.

In an individual physically dependent on opioids, administration of the recommended usual dosage of the antagonist will precipitate an acute withdrawal syndrome. The severity of the withdrawal symptoms experienced will depend on the degree of physical dependence and the dose of the antagonist administered.

If a decision is made to treat serious respiratory depression in the physically dependent patient, administration of the antagonist should be begun with care and by titration with smaller than usual doses of the antagonist.

DOSAGE AND ADMINISTRATION

Important Dosage and Administration Instructions

Pentazocine and Naloxone Tablets should be prescribed only by healthcare professionals who are knowledgeable about the use of opioids and how to mitigate the associated risks.

Use the lowest effective dosage for the shortest duration of time consistent with individual patient treatment goals [see *Warnings and Precautions*]. Reserve titration to higher doses of Pentazocine and Naloxone Tablets for patients in whom lower doses are insufficiently effective and in whom the expected benefits of using a higher dose opioid clearly outweigh the substantial risks.

Many acute pain conditions (e.g., the pain that occurs with a number of surgical procedures or acute musculoskeletal injuries) require no more than a few days of an opioid analgesic. Clinical guidelines on opioid prescribing for some acute pain conditions are available.

There is variability in the opioid analgesic dose and duration needed to adequately manage pain due both to the cause of pain and to individual patient factors. Initiate the dosing regimen for each patient individually, taking into account the patient's underlying cause and severity of pain, prior analgesic treatment and response, and risk factors for addiction, abuse, and misuse [see *Warnings*].

Respiratory depression can occur at any time during opioid therapy, especially when initiating and following dosage increases with Pentazocine and Naloxone Tablets. Consider this risk when selecting an initial dose and when making dose adjustments [see *Warnings*].

Patient Access to an Opioid Overdose Reversal Agent for the Emergency Treatment of Opioid Overdose

Inform patients and caregivers about opioid overdose reversal agents (e.g., naloxone, nalmefene). Discuss the importance of having access to an opioid overdose reversal agent, especially if the patient has risk factors for overdose (e.g., concomitant use of CNS depressants, a history of opioid use disorder, or prior opioid overdose) or if there are household members (including children) or other close contacts at risk for accidental ingestion or opioid overdose. The presence of risk factors for overdose should not prevent the management of pain in any patient [see **WARNINGS: Addiction, Abuse, and Misuse; Life-Threatening Respiratory Depression; Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**].

Discuss the options for obtaining an opioid overdose reversal agent (e.g., prescription, over-the-counter, or as part of a community-based program).

There are important differences among the opioid overdose reversal agents, such as route of administration, product strength, approved patient age range, and pharmacokinetics. Be familiar with these differences, as outlined in the approved labeling for those products, prior to recommending or prescribing such an agent.

Initial Dosage

Use of Pentazocine and Naloxone Tablets as the First Opioid Analgesic

Initiate treatment with Pentazocine and Naloxone Tablets in a dosing range of 1 tablet every 3 to 4 hours as needed for pain, at the lowest dose necessary to achieve adequate analgesia. Titrate the dose based upon the individual patient's response to their initial dose of Pentazocine and Naloxone Tablets. This may be increased to 2 tablets when needed. Total daily dosage should not exceed 12 tablets.

Conversion from Other Opioids to Pentazocine and Naloxone Tablets

There is inter-patient variability in the potency of opioid drugs and opioid formulations. Therefore, a conservative approach is advised when determining the total daily dosage of Pentazocine and Naloxone Tablets. It is safer to underestimate a patient's 24-hour Pentazocine and Naloxone Tablets dosage than to overestimate the 24-hour Pentazocine and Naloxone Tablets dosage and manage an adverse reaction due to overdose.

Titration and Maintenance of Therapy

Individually titrate Pentazocine and Naloxone Tablets to a dose that provides adequate analgesia and minimizes adverse reactions. Continually reevaluate patients receiving Pentazocine and Naloxone Tablets to assess the maintenance of pain control, signs and symptoms of opioid withdrawal, and other adverse reactions, as well as reassessing for the development of addiction, abuse, or misuse [see **WARNINGS**]. Frequent communication is important among the prescriber, other members of the healthcare team, the patient, and the caregiver/family during periods of changing analgesic requirements, including initial titration. If the level of pain increases after dosage stabilization, attempt to identify the source of increased pain before increasing the Pentazocine and Naloxone Tablets dosage. If after increasing the dosage, unacceptable opioid-related adverse reactions are observed (including an increase in pain after dosage increase), consider reducing the dosage [see *Warnings*]. Adjust the dosage to obtain an appropriate balance between management of pain and opioid-related adverse reactions.

Safe Reduction or Discontinuation of Pentazocine and Naloxone Tablets

Do not rapidly reduce or abruptly discontinue Pentazocine and Naloxone Tablets in patients who may be physically dependent on opioids. Rapid discontinuation of opioid analgesics in patients who are physically dependent on opioids has resulted in serious withdrawal symptoms, uncontrolled pain, and suicide. Rapid discontinuation has also been associated with attempts to find other sources of opioid analgesics, which may be confused with drug-seeking for abuse. Patients may also attempt to treat their pain or withdrawal symptoms with illicit opioids, such as heroin, and other substances.

When a decision has been made to decrease the dose or discontinue therapy in an opioid-dependent patient taking Pentazocine and Naloxone Tablets, there are a variety of factors that should be considered, including the total daily dose of opioid (including Pentazocine and Naloxone Tablets) the patient has been taking, the duration of treatment, the type of pain being treated, and the physical and psychological attributes of the patient. It is important to ensure ongoing care of the patient and to agree on an appropriate tapering schedule and follow-up plan so that patient and provider goals and expectations are clear and realistic. When opioid analgesics are being discontinued due to a suspected substance use disorder, evaluate and treat the patient, or refer for evaluation and treatment of the substance use disorder. Treatment should include evidence-based approaches, such as medication assisted treatment of opioid use disorder. Complex patients with co-morbid pain and substance use disorders may benefit from referral to a specialist.

There are no standard opioid tapering schedules that are suitable for all patients. Good clinical practice dictates a patient-specific plan to taper the dose of the opioid gradually. For patients on Pentazocine and Naloxone Tablets who are physically opioid-dependent, initiate the taper by a small enough increment (e.g., no greater than 10% to 25% of the total daily dose) to avoid withdrawal symptoms, and proceed with dose-lowering at an interval of every 2 to 4 weeks. Patients who have been taking opioids for briefer periods of time may tolerate a more rapid taper.

It may be necessary to provide the patient with lower dosage strengths to accomplish a successful taper. Reassess the patient frequently to manage pain and withdrawal symptoms, should they emerge. Common withdrawal symptoms include restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, myalgia, and mydriasis. Other signs and symptoms also may develop, including irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, or increased blood pressure, respiratory rate, or heart rate. If withdrawal symptoms arise, it may be necessary to pause the taper for a period of time or raise the dose of the opioid analgesic to the previous dose, and then proceed with a slower taper. In addition, evaluate patients for any changes in mood, emergence of suicidal thoughts, or use of other substances.

When managing patients taking opioid analgesics, particularly those who have been treated for an extended period of time and/or with high doses for chronic pain, ensure that a multimodal approach to pain management, including mental health support (if needed), is in place prior to initiating an opioid analgesic taper. A multimodal approach to pain management may optimize the treatment of chronic pain, as well as assist with the successful tapering of the opioid analgesic [see **WARNINGS/Withdrawal, DRUG ABUSE AND DEPENDENCE**].

HOW SUPPLIED

Pentazocine and Naloxone Tablets USP are light yellow, capsule shaped tablets debossed "NL" on left side and "680" on the right side of the bisect and plain on the other side, supplied in bottles of 100 and 500.

Bottles of 100 (NDC 43386-680-01).

Bottles of 500 (NDC 43386-680-05).

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Store Pentazocine and Naloxone Tablets securely and dispose of properly [See **PRECAUTIONS/Information for Patients**].

Dispense in a tight, light-resistant container as defined in the USP.

LUPIN and the  are registered trademarks of Lupin Pharmaceuticals, Inc.

Manufactured by:  
**Novel Laboratories, Inc.**  
Somerset, NJ 08873  
SAP Code: 281449  
Rev. 08/2025

Manufactured for:  
**Lupin Pharmaceuticals, Inc.**  
Naples, FL 34108

Medication Guide  
Pentazocine and Naloxone  
(pen taz' oh seen and nal ox' one) Tablets

Pentazocine and Naloxone Tablets are:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage moderate to severe pain, when other pain treatments such as non-opioid pain medicines do not treat your pain well enough or you cannot tolerate them.
- An opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.

Important information about Pentazocine and Naloxone Tablets:

- **Get emergency help or call 911 right away if you take too many Pentazocine and Naloxone Tablets (overdose).** When you first start taking Pentazocine and Naloxone Tablets, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur. Ask your healthcare provider about medicines like naloxone or nalmefene that can be used in an emergency to reverse an opioid overdose.
- Taking Pentazocine and Naloxone Tablets with other opioid medicines, benzodiazepines gabapentinoids (gabapentin or pregabalin), alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your Pentazocine and