33 mm

33 mm

34 mm

IMATINIB mesylate tablets, for oral use

XXXXX

IMATINIB mesylate tablets, for oral use

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use IMATINIB MESYLATE TABLETS safely and effectively. See full prescribing information for IMATINIB MESYLATE TABLETS.

IMATINIB MESYLATE tablets, for oral use
Initial U.S. Approval: 2001

----------RECENT MAJOR CHANGES----------

Warnings and Precautions, Renal Toxicity (5.14)	9/2017

----------INDICATIONS AND USAGE----------

Imatinib mesylate is a kinase inhibitor indicated for the treatment of:

- Newly diagnosed adult and pediatric patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase (1.1)
- Patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in blast crisis (BC), accelerated phase (AP), or in chronic phase (CP) after failure of interferon-alpha therapy (1.2)
- Adult patients with relapsed or refractory Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) (1.3)
- Adult patients with myelodysplastic/myeloproliferative diseases (MDS/MPD) associated with PDGFR (platelet-derived growth factor receptor) gene re-arrangements as determined with an FDA-approved test (1.5)
- Adult patients with aggressive systemic mastocytosis (ASM) without the D816V c-Kit mutation as determined with an FDA-approved test or with c-Kit mutational status unknown (1.6)
- Adult patients with hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukemia (CEL) who have the FIP1L1-PDGFRα fusion kinase (mutational analysis or FISH demonstration of CHIC2 allele deletion) and for patients with HES and/or CEL who are FIP1L1-PDGFRα fusion kinase negative or unknown (1.7)
- Adult patients with unresectable, recurrent and/or metastatic dermatofibrosarcoma protuberans (DFSP) (1.8)

----------DOSAGE AND ADMINISTRATION----------

• Adults with Ph+ CML CP (2.2):	400 mg/day
• Adults with Ph+ CML AP or BC (2.2):	600 mg/day
• Pediatrics with Ph+ CML CP (2.3):	340 mg/m²/day
• Adults with Ph+ ALL (2.4):	600 mg/day
• Adults with MDS/MPD (2.6):	400 mg/day
• Adults with ASM (2.7):	100 mg/day or 400 mg/day
• Adults with HES/CEL (2.8):	100 mg/day or 400 mg/day
• Adults with DFSP (2.9):	800 mg/day
• Patients with mild to moderate hepatic impairment (2.12):	400 mg/day
• Patients with severe hepatic impairment (2.12):	300 mg/day

All doses of Imatinib mesylate should be taken with a meal and a large glass of water. Doses of 400 mg or 600 mg should be administered once daily, whereas a dose of 800 mg should be administered as 400 mg twice a day. Imatinib mesylate can be dissolved in water or apple juice for patients having difficulty swallowing. Daily dosing of 800 mg and above should be accomplished using the 400 mg tablet to reduce exposure to iron.

----------DOSAGE FORMS AND STRENGTHS----------

Tablets (scored): 100 mg and 400 mg (3)

----------CONTRAINDICATIONS----------

None. (4)

----------WARNINGS AND PRECAUTIONS----------

- Edema and severe fluid retention have occurred. Weigh patients regularly and manage unexpected rapid weight gain by drug interruption and diuretics (5.1, 6.1)
- Cytopenias, particularly anemia, neutropenia, and thrombocytopenia, have occurred. Manage with dose reduction, dose interruption , or discontinuation of treatment. Perform complete blood counts weekly for the first month, biweekly for the second month, and periodically thereafter. (5.2)
- Severe congestive heart failure and left ventricular dysfunction have been reported, particularly in patients with comorbidities and risk factors. Monitor and treat patients with cardiac disease or risk factors for cardiac failure. (5.3)
- Severe hepatotoxicity including fatalities may occur. Assess liver function before initiation of treatment and monthly thereafter or as clinically indicated. Monitor liver function when combined with chemotherapy known to be associated with liver dysfunction. (5.4)
- Grade 3/4 hemorrhage has been reported in clinical studies in patients with newly diagnosed CML. (5.5)
- Gastrointestinal perforations, some fatal, have been reported. (5.6)
- Cardiogenic shock/left ventricular dysfunction has been associated with the initiation of Imatinib mesylate in patients with conditions associated with high eosinophil levels (e.g., HES, MDS/MPD and ASM). (5.7)
- Bullous dermatologic reactions (e.g., erythema multiforme and Stevens-Johnson syndrome) have been reported with the use of Imatinib mesylate. (5.8)
- Hypothyroidism has been reported in thyroidectomy patients undergoing levothyroxine replacement. Closely monitor TSH levels in such patients. (5.9)
- Fetal harm can occur when administered to a pregnant woman. Apprise women of the potential harm to the fetus, and to avoid pregnancy when taking Imatinib mesylate. (5.10, 8.1)
- Growth retardation occurring in children and pre-adolescents receiving Imatinib mesylate has been reported. Close monitoring of growth in children under Imatinib mesylate treatment is recommended. (5.11, 6.2)
- Tumor lysis syndrome. Close monitoring is recommended. (5.12)
- Reports of motor vehicle accidents have been received in patients receiving Imatinib mesylate. Caution patients about driving a car or operating machinery. (5.13)
- Renal toxicity. A decline in renal function may occur in patients receiving Imatinib mesylate. Evaluate renal function at baseline and during therapy, with attention to risk factors for renal dysfunction. (5.14)

----------ADVERSE REACTIONS----------

The most frequently reported adverse reactions (greater than or equal to 30%) were edema, nausea, vomiting, muscle cramps, musculoskeletal pain, diarrhea, rash, fatigue and abdominal pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Lupin Pharmaceuticals, Inc. at 1-800-399-2561 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----------DRUG INTERACTIONS----------

- CYP3A4 inducers may decrease Imatinib mesylate C_{min} and area under curve (AUC). (2.12, 7.1,12.3)
- CYP3A4 inhibitors may increase Imatinib mesylate C_{min} and AUC. (7.2,12.3)
- Imatinib mesylate is an inhibitor of CYP3A4 and CYP2D6 which may increase the C_{max} and AUC of other drugs. (7.3, 7.4, 12.3)
- Patients should inquire anticoagulation should receive low-molecular weight or standard heparin and not warfarin. (7.3)

See 17 for PATIENT COUNSELING INFORMATION

Revised: 08/2018

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE
1.1 Newly Diagnosed Philadelphia Positive Chronic Myeloid Leukemia (Ph+ CML)
1.2 Ph+ CML in Blast Crisis (BC), Accelerated Phase (AP) or Chronic Phase (CP) After Interferon-alpha (IFN) Therapy
1.3 Adult patients with Ph+ Acute Lymphoblastic Leukemia (ALL)
1.5 Myelodysplastic/Myeloproliferative Diseases (MDS/MPD)
1.6 Aggressive Systemic Mastocytosis (ASM)
1.7 Hypereosinophilic Syndrome (HES) and/or Chronic Eosinophilic Leukemia (CEL)
1.8 Dermatofibrosarcoma Protuberans (DFSP)
2 DOSAGE AND ADMINISTRATION
2.1 Drug Administration
2.2 Adult Patients with Ph+ CML CP, AP, or BC
2.3 Pediatric Patients with Ph+ CML CP
2.4 Adult Patients with Ph+ ALL
2.6 Adult Patients with MDS/MPD
2.7 Adult Patients with ASM
2.8 Adult Patients with HES/CEL
2.9 Adult Patients with DFSP
2.12 Dose Modification Guidelines
2.13 Dose Adjustment for Hepatotoxicity and Non-Hematologic Adverse Reactions
2.14 Dose Adjustment for Hematologic Adverse Reactions
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Fluid Retention and Edema
5.2 Hematologic Toxicity
5.3 Congestive Heart Failure and Left Ventricular Dysfunction
5.4 Hepatotoxicity
5.5 Hemorrhage
5.6 Gastrointestinal Disorders
5.7 Hypereosinophilic Cardiac Toxicity
5.8 Dermatologic Toxicities
5.9 Hypothyroidism
5.10 Embryo-fetal Toxicity
5.11 Growth Retardation in Children and Adolescents
5.12 Tumor Lysis Syndrome
5.13 Impairments Related to Driving and Using Machinery
5.14 Renal Toxicity
6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
6.2 Postmarketing Experience
7 DRUG INTERACTIONS
7.1 Agents Inducing CYP3A Metabolism
7.2 Agents Inhibiting CYP3A Metabolism
7.3 Interactions with Drugs Metabolized by CYP3A4
7.4 Interactions with Drugs Metabolized by CYP2D6
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.3 Females and Males of Reproductive Potential
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Hepatic Impairment
8.7 Renal Impairment
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.3 Pharmacokinetics
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
13.2 Animal Toxicology and/or Pharmacology
14 CLINICAL STUDIES
14.1 Chronic Myeloid Leukemia
14.2 Pediatric CML
14.3 Acute Lymphoblastic Leukemia
14.5 Myelodysplastic/Myeloproliferative Diseases
14.6 Aggressive Systemic Mastocytosis
14.7 Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia
14.8 Dermatofibrosarcoma Protuberans
15 REFERENCES
16 HOW SUPPLIED/STORAGE AND HANDLING
17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Newly Diagnosed Philadelphia Positive Chronic Myeloid Leukemia (Ph+ CML)
Newly diagnosed adult and pediatric patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase.

1.2 Ph+ CML in Blast Crisis (BC), Accelerated Phase (AP) or Chronic Phase (CP) After Interferon-alpha (IFN) Therapy
Patients with Philadelphia chromosome positive chronic myeloid leukemia in blast crisis, accelerated phase, or in chronic phase after failure of interferon-alpha therapy.

1.3 Adult patients with Ph+ Acute Lymphoblastic Leukemia (ALL)
Adult patients with relapsed or refractory Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL).

1.5 Myelodysplastic/Myeloproliferative Diseases (MDS/MPD)
Adult patients with myelodysplastic/myeloproliferative diseases associated with PDGFR (platelet-derived growth factor receptor) gene re-arrangements as determined with an FDA-approved test *(see Dosage and Administration (2.6))*.

1.6 Aggressive Systemic Mastocytosis (ASM)
Adult patients with aggressive systemic mastocytosis without the D816V c-Kit mutation as determined with an FDA-approved test *(see Dosage and Administration (2.7))* or with c-Kit mutational status unknown.

1.7 Hypereosinophilic Syndrome (HES) and/or Chronic Eosinophilic Leukemia (CEL)
Adult patients with hypereosinophilic syndrome and/or chronic eosinophilic leukemia who have the FIP1L1-PDGFRα fusion kinase (mutational analysis or FISH demonstration of CHIC2 allele deletion) and for patients with HES and/or CEL who are FIP1L1-PDGFRα fusion kinase negative or unknown.

1.8 Dermatofibrosarcoma Protuberans (DFSP)
Adult patients with unresectable, recurrent and/or metastatic dermatofibrosarcoma protuberans.

2 DOSAGE AND ADMINISTRATION

2.1 Drug Administration
The prescribed dose should be administered orally, with a meal and a large glass of water. Doses of 400 mg or 600 mg should be administered once daily, whereas a dose of 800 mg should be administered as 400 mg twice a day.
For patients unable to swallow the film-coated tablets, the tablets may be dispersed in a glass of water or apple juice. The required number of tablets should be placed in the appropriate volume of beverage (approximately 50 mL for a 100 mg tablet, and 200 mL for a 400 mg tablet) and stirred with a spoon. The suspension should be administered immediately after complete disintegration of the tablet(s).
For daily dosing of 800 mg and above, dosing should be accomplished using the 400 mg tablet to reduce exposure to iron.
Treatment may be continued as long as there is no evidence of progressive disease or unacceptable toxicity.

2.2 Adult Patients with Ph+ CML CP, AP, or BC
The recommended dose of Imatinib mesylate is 400 mg/day for adult patients in chronic phase CML and 600 mg/day for adult patients in accelerated phase or blast crisis.
In CML, a dose increase from 400 mg to 600 mg in adult patients with chronic phase disease, or from 600 mg to 800 mg (given as 400 mg twice daily) in adult patients in accelerated phase or blast crisis may be considered in the absence of severe adverse drug reaction and severe non-leukemia related neutropenia or thrombocytopenia in the following circumstances: disease progression (at any time), failure to achieve a satisfactory hematologic response after at least 3 months of therapy, or loss of a previously obtained hematologic response, unless it is due to cytopenia(s) not considered to be of previously achieved hematologic or cytogenetic response.

2.3 Pediatric Patients with Ph+ CML CP
The recommended dose of Imatinib mesylate for children with newly diagnosed Ph+ CML is 340 mg/m²/day (not to exceed 600 mg). Imatinib mesylate treatment can be given as a once daily dose or the daily dose may be split into two-one portion dosed in the morning and one portion in the evening. There is no experience with Imatinib mesylate treatment in children under 1 year of age.

2.4 Adults Patients with Ph+ ALL
The recommended dose of Imatinib mesylate is 600 mg/day for adult patients with relapsed/refractory Ph+ ALL.

2.6 Adult Patients with MDS/MPD
Determine PDGFRb gene rearrangements status prior to initiating treatment. Information on FDA-approved tests for the detection of PDGFRb rearrangements is available at http://www.fda.gov/companiondiagnostics.
The recommended dose of Imatinib mesylate is 400 mg/day for adult patients with MDS/MPD.

2.7 Adult Patients with ASM
Determine D816V c-Kit mutation status prior to initiating treatment. Information on FDA-approved test for the detection of D816V c-Kit mutation is available at http://www.fda.gov/companiondiagnostics.
The recommended dose of Imatinib mesylate is 400 mg/day for adult patients with ASM without the D816V c-Kit mutation. If c-Kit mutational status is not known or unavailable, treatment with Imatinib mesylate 400 mg/day may be considered for patients with ASM not responding satisfactorily to other therapies. For patients with ASM associated with eosinophilia, a clonal hematological disease related to the fusion kinase FIP1L1-PDGFRα, a starting dose of 100 mg/day is recommended. Dose increase from 100 mg to 400 mg for these patients may be considered in the absence of adverse drug reactions if assessments demonstrate an insufficient response to therapy.

2.8 Adult Patients with HES/CEL
The recommended dose of Imatinib mesylate is 400 mg/day for adult patients with HES/CEL. For HES/CEL patients with demonstrated FIP1L1-PDGFRα fusion kinase, a starting dose of 100 mg/day is recommended. Dose increase from 100 mg to 400 mg may be considered in the absence of adverse drug reactions if assessments demonstrate an insufficient response to therapy.

2.9 Adult Patients with DFSP
The recommended dose of Imatinib mesylate is 800 mg/day for adult patients with DFSP.

2.12 Dose Modification Guidelines
Concomitant Strong CYP3A4 inducers: The use of concomitant strong CYP3A4 inducers should be avoided (e.g., dexamethasone, phenytoin, carbamazepine, rifampin, rifabutin, rifampicin, phenobarbital). If patients must be coadministered a strong CYP3A4 inducer, based on pharmacokinetic studies, the dosage of Imatinib mesylate should be increased by at least 50%, and clinical response should be carefully monitored *(see Drug Interactions (7.1))*.
Hepatic impairment: Patients with mild and moderate hepatic impairment do not require a dose adjustment and should be treated per the recommended dose. A 25% decrease in the recommended dose should be used for patients with severe hepatic impairment *(see Use in Specific Populations (8.6))*.
Renal impairment: Patients with moderate renal impairment (CrCL-20 to 39 mL/min) should receive a 50% decrease in the recommended starting dose and future doses can be increased as tolerated. Doses greater than 600 mg are not recommended in patients with mild renal impairment (CrCL-40 to 59 mL/min). For patients with moderate renal impairment doses greater than 400 mg are not recommended.
Imatinib should be used with caution in patients with severe renal impairment. A dose of 100 mg/day was tolerated in two patients with severe renal impairment *(See Warnings and Precautions (5.3), Use in Specific Populations (8.7))*.

2.13 Dose Adjustment for Hepatotoxicity and Non-Hematologic Adverse Reactions
If elevations in bilirubin greater than 3 times the institutional upper limit of normal (IULN) or in liver transaminases greater than 5 times the IULN occur, Imatinib mesylate should be withheld until bilirubin levels have returned to a less than 1.5 times the IULN and transaminase levels to less than 2.5 times the IULN. In adults, treatment with Imatinib mesylate may then be continued at a reduced daily dose (i.e., 400 mg to 300 mg, 600 mg to 400 mg or 800 mg to 600 mg). In children, daily doses can be reduced under the same circumstances from 340 mg/m²/day to 260 mg/m²/day.
If a severe non-hematologic adverse reaction develops (such as severe hepatotoxicity or severe fluid retention), Imatinib mesylate should be withheld until the event has resolved. Thereafter, treatment can be resumed as appropriate depending on the initial severity of the event.

2.14 Dose Adjustment for Hematologic Adverse Reactions
Dose reduction or treatment interruptions for severe neutropenia and thrombocytopenia are recommended as indicated in Table 1.

Table 1: Dose Adjustments for Neutropenia and Thrombocytopenia

ASM associated with eosinophilia (starting dose 100 mg)	ANC[*] less than 1.0 x 10⁹/L and/or platelets less than 50 x 10⁹/L	1. Stop Imatinib mesylate until ANC greater than or equal to 1.5 x 10⁹/L and platelets greater than or equal to 75 x 10⁹/L. 2. Resume treatment with Imatinib mesylate at previous dose (i.e., dose before severe adverse reaction)
HES/CEL with FIP1L1-PDGFRα fusion kinase (starting dose 100 mg)	ANC less than 1.0 x 10⁹/L and/or platelets less than 50 x 10⁹/L	1. Stop Imatinib mesylate until ANC greater than or equal to 1.5 x 10⁹/L and platelets greater than or equal to 75 x 10⁹/L. 2. Resume treatment with Imatinib mesylate at previous dose (i.e., dose before severe adverse reaction)
Chronic Phase CML (starting dose 400 mg) MDS/MPD, ASM and HES/CEL (starting dose 400 mg)	ANC less than 1.0 x 10⁹/L and/or platelets less than 50 x 10⁹/L	1. Stop Imatinib mesylate until ANC greater than or equal to 1.5 x 10⁹/L and platelets greater than or equal to 75 x 10⁹/L. 2. Resume treatment with Imatinib mesylate at the original starting dose of 400 mg. 3. If recurrence of ANC less than 1.0 x 10⁹/L and/or platelets less than 50 x 10⁹/L, repeat step 1 and resume Imatinib mesylate at a reduced dose of 300 mg.
Ph+ CML: Accelerated Phase and Blast Crisis (starting dose 600 mg) Ph+ ALL (starting dose 600 mg)	ANC less than 0.5 x 10⁹/L and/or platelets less than 10 x 10⁹/L	1. Check if cytopenia is related to leukemia (marrow aspirate or biopsy) 2. If cytopenia is unrelated to leukemia, reduce dose of Imatinib mesylate to 400 mg. 3. If cytopenia persists 2 weeks, reduce further to 300 mg. 4. If cytopenia persists 4 weeks and is still unrelated to leukemia, stop Imatinib mesylate until ANC greater than or equal to 1 x 10⁹/L and platelets greater than or equal to 20 x 10⁹/L, and then resume treatment at 300 mg.
DFSP (starting dose 800 mg)	ANC less than 1.0 x 10⁹/L and/or platelets less than 50 x 10⁹/L	1. Stop Imatinib mesylate until ANC greater than or equal to 1.5 x 10⁹/L and platelets greater than or equal to 75 x 10⁹/L. 2. Resume treatment with Imatinib mesylate at dose of 600 mg. 3. In the event of recurrence of ANC less than 1.0 x 10⁹/L and/or platelets less than 50 x 10⁹/L, repeat step 1 and resume Imatinib mesylate at reduced dose of 400 mg.
Pediatric newly diagnosed chronic phase CML (starting dose 340 mg/m²)	ANC less than 1.0 x 10⁹/L and/or platelets less than 50 x 10⁹/L	1. Stop Imatinib mesylate until ANC greater than or equal to 1.5 x 10⁹/L and platelets greater than or equal to 75 x 10⁹/L. 2. Resume treatment with Imatinib mesylate at previous dose (i.e., dose before severe adverse reaction). 3. In the event of recurrence of ANC less than 1.0 x 10⁹/L and/or platelets less than 50 x 10⁹/L, repeat step 1 and resume Imatinib mesylate at reduced dose of 260 mg/m².

[*] ANC = absolute neutrophil count

3 DOSAGE FORMS AND STRENGTHS
100 mg film-coated tablets
Brownish, round, convex, film-coated tablets debossed with "100" on one side and score line on other side with "N" on one side of the score line and "I" on other side of score line.
400 mg film-coated tablets
Brownish, oval, biconvex, film-coated tablets debossed with "400" on one side and score line on other side with "N" on one side of the score line and "I" on other side of score line.

4 CONTRAINDICATIONS
None.

5 WARNINGS AND PRECAUTIONS

5.1 Fluid Retention and Edema
Imatinib mesylate is often associated with edema and occasionally serious fluid retention *(see Adverse Reactions (6.1))*. Weigh and monitor patients regularly for signs and symptoms of fluid retention. Investigate unexpected rapid weight gain carefully and provide appropriate treatment. The probability of edema was increased with higher Imatinib mesylate dose and age greater than 65 years in the CML studies. Severe superficial edema was reported in 1.5% of newly diagnosed CML patients taking Imatinib mesylate, and in 2% to 6% of other adult CML patients taking Imatinib mesylate. In addition, other severe fluid retention (e.g., pleural effusion, pericardial effusion, pulmonary edema, and ascites) reactions were reported in 1.3% of newly diagnosed CML patients taking Imatinib mesylate, and in 2% to 6% of other adult CML patients taking Imatinib mesylate. In a randomized trial in patients with newly diagnosed Ph+CML in chronic phase comparing Imatinib mesylate and nilotinib, severe (Grade 3 or 4) fluid retention occurred in 2.5% of patients receiving Imatinib mesylate and in 3.9% of patients receiving nilotinib 300 mg twice daily. Effusions (including pleural effusion, pericardial effusion, ascites) or pulmonary edema were observed in 2.1% (none were Grade 3 or 4) of patients in the Imatinib mesylate arm and 2.2% (0.7% Grade 3 or 4) of patients in the nilotinib 300 mg twice daily arm.

5.2 Hematologic Toxicity
Treatment with Imatinib mesylate is associated with anemia, neutropenia, and thrombocytopenia. Perform complete blood counts weekly for the first month, biweekly for the second month, and periodically thereafter as clinically indicated (for example, every 2 to 3 months). In CML, the occurrence of these cytopenias is dependent on the stage of disease and is more frequent in patients with accelerated phase CML or blast crisis than in patients with chronic phase CML. In pediatric CML patients the most frequent toxicities observed were Grade 3 or 4 cytopenias including neutropenia, thrombocytopenia and anemia. These generally occur within the first several months of therapy *(see Dosage and Administration (2.14))*.

5.3 Congestive Heart Failure and Left Ventricular Dysfunction
Congestive heart failure and left ventricular dysfunction have been reported in patients taking Imatinib mesylate. Cardiac adverse reactions were more frequent in patients with advanced age or co-morbidities including previous medical history of cardiac disease. In an international randomized Phase 3 study in 1,106 patients with newly diagnosed Ph+ CML in chronic phase, severe cardiac failure and left ventricular dysfunction were observed in 0.7% of patients taking Imatinib mesylate compared to 0.9% of patients taking IFN + Ara-C. In another randomized trial with newly diagnosed Ph+ CML patients in chronic phase that compared Imatinib mesylate and nilotinib, cardiac failure was observed in 1.1% of patient in the Imatinib mesylate arm and 2.2% of patients in the nilotinib 300 mg twice daily arm and severe (Grade 3 or 4) cardiac failure occurred in 0.7% of patients in each group. Carefully monitor patients with cardiac disease or risk factors for cardiac or history of renal failure. Evaluate and treat any patient with signs or symptoms consistent with cardiac or renal failure.

5.4 Hepatotoxicity
Hepatotoxicity, occasionally severe, may occur with Imatinib mesylate *(see Adverse Reactions (6.1))*. Cases of total liver failure and severe liver injury requiring liver transplants have been reported with both short-term and long-term use of Imatinib mesylate. Monitor liver function (transaminases, bilirubin, and alkaline phosphatase) before initiation of treatment and monthly, or as clinically indicated. Manage laboratory abnormalities with Imatinib mesylate interruption and/or dose reduction *(see Dosage and Administration (2.13))*. When Imatinib mesylate is combined with chemotherapy, liver toxicity in the form of transaminase elevation and hyperbilirubinemia has been observed. Additionally, there have been reports of acute liver failure. Monitoring of hepatic function is recommended.

5.5 Hemorrhage
In a trial of Imatinib mesylate versus IFN+Ara-C in patients with the newly diagnosed CML, 1.8% of patients had Grade 3/4 hemorrhage. In a randomized trial in patients with newly diagnosed Ph+ CML in chronic phase comparing Imatinib mesylate and nilotinib, GI hemorrhage occurred in 1.4% of patients in the Imatinib mesylate arm, and in 2.9% of patients in the nilotinib 300 mg twice daily arm. None of these events were Grade 3 or 4 in the Imatinib mesylate arm; 0.7% were Grade 3 or 4 in the nilotinib 300 mg twice daily arm. In addition, gastric antral vascular ectasia has been reported in postmarketing experience.

5.6 Gastrointestinal Disorders
Imatinib mesylate is sometimes associated with GI irritation. Imatinib mesylate should be taken with food and a large glass of water to minimize this problem. There have been rare reports, including fatalities, of gastrointestinal perforation.

5.7 Hypereosinophilic Cardiac Toxicity
In patients with hypereosinophilic syndrome with occult infiltration of HES cells within the myocardium, cases of cardiogenic shock/left ventricular dysfunction have been associated with HES cell degranulation upon the initiation of Imatinib mesylate therapy. The condition was reported to be reversible with the administration of systemic steroids, circulatory support measures and temporary suspension of Imatinib mesylate treatment.
Myelodysplastic/myeloproliferative disease and systemic mastocytosis may be associated with high eosinophil levels. Consider performing an echocardiogram and determining serum tryptase in patients with HES/CEL, and in patients with MDS/MPD or ASM associated with high eosinophil levels. If either is abnormal, consider prophylactic use of systemic steroids (1 to 2 mg/kg) for one to two weeks concurrently with Imatinib mesylate at the initiation of therapy.

5.8 Dermatologic Toxicities
Bullous dermatologic reactions, including erythema multiforme and Stevens-Johnson syndrome, have been reported with use of Imatinib mesylate. In some cases of bullous dermatologic reactions, including erythema multiforme and Stevens-Johnson syndrome, symptoms reported were preceded by fever and flu-like symptoms; a recurrent dermatologic reaction was observed upon rechallenge. Several foreign postmarketing reports have described cases in which patients tolerated the reintroduction of Imatinib mesylate therapy after resolution or improvement of the bullous reaction. In these instances, Imatinib mesylate was resumed at a dose lower than that at which the reaction occurred and some patients also received concomitant treatment with corticosteroids or antihistamines.

5.9 Hypothyroidism
Clinical cases of hypothyroidism have been reported in thyroidectomy patients undergoing levothyroxine replacement during treatment with Imatinib mesylate. Closely monitor TSH levels in such patients.

5.10 Embryo-fetal Toxicity
Imatinib mesylate can cause fetal harm when administered to a pregnant woman. Imatinib mesylate was teratogenic in rats when administered during organogenesis at doses approximately equal to the maximum human dose of 800 mg/day based on body surface area. Significant post-implantation loss was seen in female rats administered imatinib mesylate at doses approximately one-half the maximum human dose of 800 mg/day based on body surface area. Advise sexually active female patients of reproductive potential to use effective contraception (methods that result in less than 1% pregnancy rates) when administering Imatinib mesylate and for 14 days after stopping Imatinib mesylate. If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, apprise the patient of the potential hazard to a fetus *(see Use in Specific Populations (8.1))*.

5.11 Growth Retardation in Children and Adolescents
Growth retardation has been reported in children and pre-adolescents receiving Imatinib mesylate. The long term effects of prolonged treatment with Imatinib mesylate on growth in children are unknown. Therefore, monitor growth in children under Imatinib mesylate treatment *(see Adverse Reactions (6.1))*.

5.12 Tumor Lysis Syndrome
Cases of Tumor Lysis Syndrome (TLS), including fatal cases, have been reported in patients with CML, ALL and myelodysplastic syndrome receiving Imatinib mesylate. The patients at risk of TLS are those with high tumor burden, having a high proliferative rate or high tumor burden prior to treatment. Monitor these patients closely and take appropriate precautions. Due to the possible occurrence of TLS, correct clinically significant dehydration and treat high uric acid levels prior to initiation of Imatinib mesylate.

5.13 Impairments Related to Driving and Using Machinery
Motor vehicle accidents have been reported in patients receiving Imatinib mesylate. Advise patients that they may experience side effects such as dizziness, blurred vision or somnolence during treatment with Imatinib mesylate. Recommend caution when driving a car or operating machinery.

5.14 Renal Toxicity
A decline in renal function may occur in patients receiving Imatinib mesylate. Median estimated glomerular filtration rate (eGFR) values in patients on imatinib mesylate 400 mg daily for newly-diagnosed CML (four randomized trials) and another clinical trial declined from a baseline value of 95 mL/min/1.73m² (N=1190) to 75 mL/min/1.73m² at 12 months (N=1082) and 69 mL/min/1.73m² at 60 months (N=549). Evaluate renal function prior to initiating Imatinib mesylate and monitor during therapy, with attention to risk factors for renal dysfunction such as pre-existing renal impairment, diabetes mellitus, hypertension, and congestive heart failure.

6 ADVERSE REACTIONS
The following serious adverse reactions are described elsewhere in the labeling:
- Fluid Retention and Edema *[see Warnings and Precautions (5.1)]*
- Hematologic Toxicity *[see Warnings and Precautions (5.2)]*
- Congestive Heart Failure and Left Ventricular Dysfunction *[see Warnings and Precautions (5.3)]*
- Hepatotoxicity *[see Warnings and Precautions (5.4)]*
- Hemorrhage *[see Warnings and Precautions (5.5)]*
- Gastrointestinal Disorders *[see Warnings and Precautions (5.6)]*
- Hypereosinophilic Cardiac Toxicity *[see Warnings and Precautions (5.7)]*
- Dermatologic Toxicities *[see Warnings and Precautions (5.8)]*
- Hypothyroidism *[see Warnings and Precautions (5.9)]*
- Growth Retardation in Children and Adolescents *[see Warnings and Precautions (5.11)]*
- Tumor Lysis Syndrome *[see Warnings and Precautions (5.12)]*
- Impairments Related to Driving and Using Machinery *[see Warnings and Precautions (5.13)]*
- Renal Toxicity *[see Warnings and Precautions (5.14)]*

6.1 Clinical Trials Experience
Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Chronic Myeloid Leukemia
The majority of Imatinib mesylate-treated patients experienced adverse reactions at some time. Imatinib mesylate was discontinued due to drug-related adverse reactions in 2.4% of patients receiving Imatinib mesylate in the randomized trial of newly diagnosed patients after IFN failure, in 4.3% of patients receiving Imatinib mesylate in the randomized trial of newly diagnosed patients with Ph+ CML in chronic phase compared to Imatinib mesylate and nilotinib, and in 12.5% of patients receiving Imatinib mesylate in the randomized trial of newly diagnosed patients with Ph+ CML in chronic phase compared to Imatinib mesylate and dasatinib. Imatinib mesylate dose was discontinued due to drug-related adverse reactions in 4% of patients in chronic phase after failure of interferon-alpha therapy, in 4% of patients in accelerated phase and in 5% of patients in blast crisis.
The most frequently reported drug-related adverse reactions were edema, nausea and vomiting, muscle cramps, musculoskeletal pain, diarrhea, rash (Table 2 and Table 3 for newly diagnosed CML and Table 4 for other patients). Edema was most frequently periorbital or in lower limbs and was managed with diuretics, other supportive measures, or by dose reduction of Imatinib mesylate *(see Dosage and Administration (2.13))*. The frequency of severe superficial edema was 1.5% to 6%.
A variety of adverse reactions represent local or general fluid retention including pleural effusion, ascites, pulmonary edema and rapid weight gain with or without dyspnea. These reactions appear to be dose related, were most common in the blast crisis and accelerated phase studies (where the dose was 600 mg/day), and are more common in the elderly. These reactions may require interruption of Imatinib mesylate treatment and using diuretics or other appropriate supportive care measures. These reactions may be serious or life threatening.
Adverse reactions, regardless of relationship to study drug, that were reported in at least 10% of the Imatinib mesylate treated patients are shown in Tables 2, 3 and 4.

Table 2: Adverse Reaction Regardless of Relationship to Study Drug Reported in Newly Diagnosed CML Clinical Trial in the Imatinib mesylate versus IFN+Ara-C Study (greater than or equal to 10% of Imatinib mesylate Treated Patients[1])

	All Grades		CTC Grades 3/4	
Preferred Term	Imatinib mesylate N=551 (%)	IFN+Ara-C N=533 (%)	Imatinib mesylate N=551 (%)	IFN+Ara-C N=533 (%)
Fluid Retention	61.7	11.1	2.5	0.9
– Superficial Edema	58.9	9.6	1.5	0.4
– Other Fluid Retention Reactions[2]	6.9	1.9	1.3	0.6
Nausea	49.5	61.5	1.3	5.1
Muscle Cramps	49.2	11.8	2.2	0.2
Musculoskeletal Pain	47.0	44.8	5.4	8.6
Diarrhea	45.4	43.3	3.3	3.2
Rash and Related Terms	40.1	26.1	2.9	2.4
Fatigue	38.8	67.0	1.8	25.1
Headache	37.0	43.3	0.5	3.8
Joint Pain	31.4	38.1	2.5	7.7
Abdominal Pain	36.5	25.9	4.2	3.9
Nasopharyngitis	30.5	8.8	0	0.4
Hemorrhage	28.9	21.2	1.8	1.7
– GI Hemorrhage	1.6	1.1	0.5	0.2
– CNS Hemorrhage	0.2	0.4	0	0.4
Myalgia	24.1	38.8	1.5	8.3
Dyspepsia	22.5	27.8	2.0	3.4
Dyspepsia	18.9	8.3	0	0.8
Cough	20.0	23.1	0.2	0.6
Pharyngolaryngeal Pain	18.1	11.4	0.2	0
Upper Respiratory Tract Infection	21.2	8.4	0.2	0.4
Dizziness	19.4	24.4	0.9	3.8
Pyrexia	17.8	42.6	0.9	3.0
Weight Increased	15.6	2.6	2.0	0.4
Insomnia	14.7	18.6	0	2.3
Depression	14.9	35.8	0.5	13.1
Influenza	13.8	6.2	0.2	0.2
Bone Pain	11.3	15.6	1.6	3.4
Constipation	11.4	14.4	0.7	0.2
Sinusitis	11.4	6.0	0.2	0.2

[1] All adverse reactions occurring in greater than or equal to 10% of Imatinib mesylate treated patients listed regardless of suspected regarding relationship to treatment.
[2] Other fluid retention reactions include pleural effusion, ascites, pulmonary edema, pericardial effusion, anasarca, edema aggravated, and fluid retention not otherwise specified.

**Table 3: Most Frequently Reported Non-hematologic Adverse Reactions (Regardless of Relationship to Study Dr

Musculoskeletal and Connective Tissue Disorders: avascular necrosis/hip osteonecrosis, rhabdomyolysis/myopathy, growth retardation in children, musculoskeletal pain upon treatment discontinuation (including myalgia, pain in extremity, arthralgia, bone pain)

Reproductive Disorders: hemorrhagic corpus luteum/hemorrhagic ovarian cyst

Blood and Lymphatic System Disorders: thrombotic microangiopathy

Including some fatalities

7 DRUG INTERACTIONS

7.1 Agents Inducing CYP3A Metabolism

Concomitant administration of Imatinib mesylate and strong CYP3A4 inducers may reduce total exposure of imatinib; consider alternative agents *[see Clinical Pharmacology (12.3)]*.

7.2 Agents Inhibiting CYP3A Metabolism

Concomitant administration of Imatinib mesylate and strong CYP3A4 inhibitors may result in a significant imatinib exposure increase. Grapefruit juice may also increase plasma concentrations of imatinib; avoid grapefruit juice *[see Clinical Pharmacology (12.3)]*.

7.3 Interactions with Drugs Metabolized by CYP3A4

Imatinib mesylate will increase plasma concentration of CYP3A4 metabolized drugs (e.g., triazolo-benzodiazepines, dihydropyridine calcium channel blockers, certain HMG-CoA reductase inhibitors, etc.). Use caution when administering imatinib mesylate with these CYP3A4 substrates that have a narrow therapeutic window.

Because warfarin is metabolized by CYP2C9 and CYP3A4, use low-molecular weight or standard heparin instead of warfarin in patients who require anticoagulation *[see Clinical Pharmacology (12.3)]*.

7.4 Interactions with Drugs Metabolized by CYP2D6

Use caution when administering Imatinib mesylate with CYP2D6 substrates that have a narrow therapeutic window.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Imatinib mesylate can cause fetal harm when administered to a pregnant woman based on human and animal data. There are no clinical studies regarding use of Imatinib mesylate in pregnant women. There have been postmarket reports of spontaneous abortions and congenital anomalies from women who were exposed to Imatinib mesylate during pregnancy. Reproductive studies in rats have demonstrated that imatinib mesylate induced teratogenicity and increased incidence of congenital abnormalities and embryofetal lethality at exposures (approximately equal to the maximum human dose of 800 mg/day based on body surface area), the number of fetuses with encephalocoele and exencephaly was higher than historical control values and these findings were associated with missing or underdeveloped cranial bones. Lower mean fetal body weights were associated with retarded skeletal ossification.

In rabbits, at doses 1.5 times higher than the maximum human dose of 800 mg/day based on body surface area, no effects on the reproductive parameters with regard to implantation sites, number of live fetuses, sex ratio or fetal weight were observed. The examinations of the fetuses did not reveal any drug related morphological changes.

In a pre- and postnatal development study in rats, pregnant rats received oral doses of imatinib mesylate during gestation (organogenesis) and lactation up to 45 mg/kg/day. Five animals developed a red vaginal discharge in the 45 mg/kg/ day group on Days 14 or 15 of gestation, the significance of which is unknown since all females produced viable litters and none had increased post-implantation loss. Other maternal effects noted only at the dose of 45 mg/kg/day (approximately one-half the maximum human dose of 800 mg/day based on body surface area) included an increased numbers of stillborn pups and pups dying between postpartum Days 0 and 4. In the F, offspring at this same dose level, mean body weights were reduced from birth until terminal sacrifice and the number of litters achieving criterion for preputial separation was slightly decreased. There were no other significant effects in developmental parameters or behavioral testing. F, fertility was not affected but reproductive effects were noted at 45 mg/kg/day including an increased number of resorptions and a decreased number of viable fetuses. The NOEL for both maternal animals and the F, generation was 15 mg/kg/day.

8.2 Lactation

Risk summary

Imatinib and its active metabolite are excreted into human milk. Because of the potential for serious adverse reactions in breastfed infants from Imatinib mesylate, advise a lactating woman not to breastfeed during treatment and for 1 month after the last dose.

Human Data

Based on data from 3 breastfeeding women taking Imatinib mesylate, the milk-plasma ratio is about 0.5 for imatinib and about 0.9 for the active metabolite. Considering the combined concentration of imatinib and active metabolite, a breastfed infant could receive up to 10% of the maternal therapeutic dose based on body weight.

8.3 Females and Males of Reproductive Potential

Pregnancy Testing

Human postmarketing reports and animal studies have shown Imatinib mesylate to be harmful to the developing fetus. Test pregnancy status in females with reproductive potential prior to the initiation of treatment with Imatinib mesylate.

Contraception

Females

Advise female patients of reproductive potential to use effective contraception (methods that result in less than 1 % pregnancy rates) when using Imatinib mesylate during treatment and for fourteen days after stopping treatment with Imatinib mesylate *[see Use in Specific Populations (8.1)]*.

Infertility

The risk of infertility in females or males of reproductive potential has not been studied in humans. In a rat study, the fertility in males and females was not affected *[see Nonclinical Toxicology (13)]*.

8.4 Pediatric Use

The safety and effectiveness of Imatinib mesylate have been demonstrated in pediatric patients with newly diagnosed Ph+ chronic phase CML *[see Clinical Studies (14.2)]*. There are no data in children under 1 year of age.

8.5 Geriatric Use

In the CML clinical studies, approximately 20% of patients were older than 65 years. In the study of patients with newly diagnosed CML, 6% of patients were older than 65 years. The frequency of edema was higher in patients older than 65 years as compared to younger patients; no other difference in the safety profile was observed *[see Warnings and Precautions (5.1)]*. The clinical study of Imatinib mesylate was similar in older and younger patients.

8.6 Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of both imatinib and its major metabolite, CGP74588, was assessed in 84 patients with cancer with varying degrees of hepatic impairment at imatinib doses ranging from 100 mg to 800 mg.

Mild and moderate hepatic impairment do not influence exposure to imatinib and CGP74588. In patients with severe hepatic impairment (bilirubin levels greater than 3 times ULN), imatinib C_{max} and area under curve (AUC) increased by 63% and 45% and the CGP74588 C_{max} and AUC increased by 56% and 55%, relative to patients with normal hepatic function *[see Clinical Pharmacology (12.2)]*. Reduce the dose by 25% for patients with severe hepatic impairment *[see Dosage and Administration (2.12)]*.

Table 16: Liver Function Classification

Liver Function Test	Normal (n=18)	Mild (n=38)	Moderate (n=28)	Severe (n=26)
Total Bilirubin	less than or equal to ULN	greater than 1.0 to 1.5 times the ULN	greater than 1.5 to 3 times the ULN	greater than 3 to 10 times the ULN
SGOT	less than or equal to ULN	greater than ULN can be normal if Total Bilirubin is greater than ULN	Any	Any

ULN=upper limit of normal for the institution

8.7 Renal Impairment

The effect of renal impairment on the pharmacokinetics of imatinib was assessed in 59 patients with cancer and varying degrees of renal impairment at single and steady state imatinib doses ranging from 100 to 800 mg/day. The mean exposure to imatinib (dose normalized AUC) in patients with mild and moderate renal impairment increased 1.5- to 2-fold compared to patients with normal renal function. Patients with severe renal impairment *[see Clinical Pharmacology (12.3)]*. Dose reductions are necessary for patients with moderate and severe renal impairment *[see Dosage and Administration (2.12)]*.

Table 17: Renal Function Classification

Renal Dysfunction	Renal Function Tests
Mild	CrCL = 40 to 59 mL/min
Moderate	CrCL = 20 to 39 mL/min
Severe	CrCL = less than 20 mL/min

CrCL = Creatinine Clearance

10 OVERDOSAGE

Experience with doses greater than 800 mg is limited. Isolated cases of Imatinib mesylate overdose have been reported. In the event of overdosage, observe the patient and give appropriate supportive treatment.

Adult Overdose

1,200 to 1,600 mg (duration varying between 1 to 10 days): Nausea, vomiting, diarrhea, rash erythema, edema, swelling, fatigue, muscle spasms, thrombocytopenia, pancytopenia, abdominal pain, headache, decreased appetite.

1,800 to 3,200 mg (as high as 3,200 mg daily for 6 days): Weakness, myalgia, increased CPK, increased bilirubin, gastrointestinal pain.

6,400 mg (single dose): One case in the literature reported one patient who experienced nausea, vomiting, abdominal pain, pyrexia, facial swelling, neutrophil count decreased, increase transaminases.

8 to 10 g (single dose): Vomiting and gastrointestinal pain have been reported.

A patient with myeloid blast crisis experienced Grade 1 elevations of serum creatinine, Grade 2 ascites and elevated liver transaminase levels, and Grade 3 elevations of bilirubin after inadvertently taking 1,200 mg of Imatinib mesylate daily for 6 days. Therapy was temporarily interrupted and complete reversal of all abnormalities occurred within 1 week. Treatment was resumed at a dose of 400 mg daily without recurrence of adverse reactions. Another patient developed severe muscle cramps after taking 1,600 mg of Imatinib mesylate daily for 6 days. Complete resolution of muscle cramps occurred following interruption of therapy and treatment was subsequently resumed. Another patient that was prescribed 400 mg daily, took 800 mg of Imatinib mesylate on Day 1 and 1,200 mg on Day 2. Therapy was interrupted, no adverse reactions occurred and the patient resumed therapy.

Pediatric Overdose

One 3-year-old male exposed to a single dose of 400 mg experienced vomiting, diarrhea and anorexia and another 3-year-old male exposed to a single dose of 980 mg experienced decreased white blood cell count and diarrhea.

11 DESCRIPTION

Imatinib is a small molecule kinase inhibitor. Imatinib mesylate film-coated tablets contain imatinib mesylate equivalent to 100 mg or 400 mg of imatinib free base. Imatinib mesylate is designated chemically as 4-((4-Methylpiperazin-1-yl) methyl)-N-(4-methyl-3-((4-(pyridin-3-yl)pyrimidin-2-yl)amino)phenyl)benzamide methanesulfonate and its structural formula is:



Imatinib mesylate is a white to off-white crystalline powder. Its molecular formula is $C_{23}H_{31}N_7O \cdot CH_4SO_3H$ and its molecular weight is 589.72. Imatinib mesylate is freely soluble in water. Sparingly soluble in Methanol. Slightly soluble in ethanol (96 percent), *practically insoluble in methylene chloride.*

Inactive Ingredients: Opadry brown (03A565002), purified water and sodium stearyl fumarate.

Opadry brown (03A565002) contains hypromellose, iron oxide red, iron oxide yellow, and talc.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Imatinib mesylate is a protein-tyrosine kinase inhibitor that inhibits the BCR-ABL tyrosine kinase, the constitutive abnormal tyrosine kinase created by the Philadelphia chromosome abnormality in CML. Imatinib inhibits proliferation and induces apoptosis in BCR-ABL positive cell lines as well as fresh leukemic cells from Philadelphia chromosome positive chronic myeloid leukemia. Imatinib inhibits colony formation in assays using *ex vivo* peripheral blood and bone marrow samples from CML patients.

In vivo, imatinib inhibits tumor growth of BCR-ABL transfected murine myeloid cells as well as BCR-ABL positive leukemia lines derived from CML patients in blast crisis.

Imatinib is also an inhibitor of the receptor tyrosine kinases for platelet-derived growth factor (PDGF) and stem cell factor (SCF), c-kit, and inhibits PDGF- and SCF-mediated cellular events.

12.3 Pharmacokinetics

The pharmacokinetics of Imatinib mesylate have been evaluated in healthy subjects and in population pharmacokinetic studies in over 900 patients.

Absorption and Distribution

Imatinib is well absorbed after oral administration with C_{max} achieved within 2 to 4 hours post-dose. Mean absolute bioavailability is 98%. Mean imatinib AUC increases proportionally with increasing doses ranging from 25 mg to 1,000 mg. There is no significant change in the pharmacokinetics of imatinib on repeated dosing, and accumulation is 1.5- to 2.5-fold at steady state when Imatinib mesylate is dosed once-daily. At clinically relevant concentrations of imatinib, binding to plasma proteins in *in vitro* experiments is approximately 95%, mostly to albumin and α1-acid glycoprotein.

Elimination

Metabolism

CYP3A4 is the major enzyme responsible for metabolism of imatinib. Other cytochrome P450 enzymes, such as CYP1A2, CYP2C9, CYP2C19, and CPY21B, play a minor role in its metabolism. The main circulating active metabolite in humans is the N-demethylated piperazine derivative, formed predominantly by CYP3A4. It shows *in vitro* potency similar to the parent imatinib. The plasma AUC for this metabolite is about 15% of the AUC for imatinib. The plasma protein binding of N-demethylated metabolite CGP74588 is similar to that of the parent compound. Human liver microsome studies demonstrated that Imatinib mesylate is a potent competitive inhibitor of CYP2C9, CYP2D6, and CYP3A4/5 with Ki values of 27, 7.5, and 8 µM, respectively.

Excretion

Imatinib elimination is predominately in the feces, mostly as metabolites. Based on the recovery of compound(s) after an oral ^{14}C-labeled dose of imatinib, approximately 81% of the dose was eliminated within 7 days, in feces (68% of dose) and urine (13% of dose). Unchanged imatinib accounted for 25% of the dose (5% urine, 20% feces), the remainder being metabolites.

Following oral administration in healthy volunteers, the elimination half-lives of imatinib and its major active metabolite, the N-demethylated derivative (CGP74588), are approximately 18 and 40 hours, respectively.

Typically, clearance of imatinib in a 50-year-old patient weighing 50 kg is expected to be 8 L/h, while for a 50-year-old patient weighing 100 kg the clearance will increase to 14 L/h. The inter-patient variability of 40% in clearance does not warrant initial dose adjustment based on body weight and/or age but indicates the need for close monitoring for treatment-related toxicity.

Specific Populations

Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of both imatinib and its major metabolite, CGP74588, was assessed in 84 patients with cancer and varying degrees of hepatic impairment *[see Use in Specific Populations (8.6)]* at imatinib doses ranging from 100 mg to 800 mg. Exposure to both imatinib and CGP74588 was comparable between each of the mildly and moderately hepatically-impaired groups and the normal group. Patients with severe hepatic impairment tend to have lower plasma exposure to both imatinib and its metabolite than patients with normal hepatic function. At steady state, the mean C_{max} dose and AUC dose for imatinib increased by about 63% and 45%, respectively, in patients with severe hepatic impairment compared to patients with normal hepatic function. The mean C_{max} dose and AUC dose for CGP74588 increased by about 56% and 55%, respectively, in patients with severe hepatic impairment compared to patients with normal hepatic function. Dose reductions are necessary for patients with severe hepatic impairment *[see Dosage and Administration (2.12)]*.

Renal Impairment

The effect of renal impairment on the pharmacokinetics of imatinib was assessed in 59 cancer patients with varying degrees of renal impairment *[see Use in Specific Populations (8.7)]* at single and steady state imatinib doses ranging from 100 to 800 mg/day. The mean exposure to imatinib (dose normalized AUC) in patients with mild and moderate renal impairment increased 1.5- to 2-fold compared to patients with normal renal function. The AUCs did not increase for dosage regimens higher than 600 mg in patients with mild renal impairment. The AUCs did not increase for dosage regimens higher than 400 mg in patients with moderate renal impairment. Two patients with severe renal impairment were dosed with 100 mg/day and their exposures were found to be similar to those seen in patients with normal renal function receiving 400 mg/day. Dose reductions are necessary for patients with moderate and severe renal impairment *[see Dosage and Administration (2.12)]*.

Pediatric Use

As in adult patients, imatinib was rapidly absorbed after oral administration in pediatric patients, with a C_{max} of 2 to 4 hours. Apparent oral clearance was similar to adult values (11.0 L/h/m² in children vs. 10.0 L/h/m² in adults), as was the half-life (14.8 hours in children vs. 17.1 hours in adults). Dosing in children at both 260 mg/m² and 340 mg/m² achieved an AUC similar to the 400 mg dose in adults. The comparison of AUC on Day 8 vs. Day 1 at 260 mg/m² and 340 mg/m² dose levels revealed a 1.5- and 2.2-fold drug accumulation, respectively, after repeated once-daily dosing. Mean imatinib AUC did not increase proportionally with increasing dose.

Based on population pharmacokinetic analysis in pediatric patients with hematological disorders (CML or other hematological disorders treated with imatinib), clearance of imatinib increases with increasing body surface area (BSA). After correcting for the BSA effect, other demographics such as age, body weight and body mass index did not have clinically significant effects on the exposure of imatinib. The analysis confirmed that exposure of imatinib in pediatric patients receiving 260 mg/m² once-daily (not exceeding 400 mg once-daily) or 340 mg/m² once-daily (not exceeding 600 mg once-daily) were similar to those in adult patients who received imatinib 400 mg or 600 mg once-daily.

Drug Interactions

Agents Inducing CYP3A Metabolism

Pretreatment of healthy volunteers with multiple doses of rifampin followed by a single dose of imatinib mesylate, increased imatinib mesylate oral-dose clearance by 3.8-fold, which significantly (p less than 0.05) decreased mean C_{max} and AUC.

Similar findings were observed in patients receiving 400 to 1200 mg/day imatinib mesylate concomitantly with enzyme-inducing anti-epileptic drugs (EIAED) (e.g., carbamazepine, oxcarbamazepine, phenytoin, fosphenytoin, phenobarbital, and primidone). The mean dose normalized AUC for imatinib in the patients receiving EIAED's decreased by 73% compared to patients not receiving EIAED.

Concomitant administration of Imatinib mesylate and St. John's Wort led to a 30% reduction in the AUC of imatinib.

Consider alternative therapeutic agents with less enzyme induction potential in patients when rifampin or other CYP3A4 inducers are indicated. Imatinib mesylate doses up to 1200 mg/day (600 mg twice daily) have been given to patients receiving concomitant strong CYP3A4 inducers *[see Dosage and Administration (2.12)]*.

Agents Inhibiting CYP3A Metabolism

There was a significant increase in exposure to imatinib (mean C_{max} and AUC increased by 26% and 40%, respectively) in healthy subjects when Imatinib mesylate was coadministered with a single dose of ketoconazole (a CYP3A4 inhibitor). Caution is recommended when administering Imatinib mesylate with strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, clarithromycin, atazanavir, indinavir, nefazodone, nelfinavir, ritonavir, saquinavir, telithromycin, and voriconazole). Grapefruit juice may also increase plasma concentrations of imatinib and should be avoided.

Interactions with Drugs Metabolized by CYP3A4

Imatinib mesylate increases the mean C_{max} and AUC of simvastatin (CYP3A4 substrate) 2- and 3.5-fold, respectively, suggesting an inhibition of the CYP3A4 by Imatinib mesylate. Particular caution is recommended when administering Imatinib mesylate with CYP3A4 substrates that have a narrow therapeutic window (e.g., alfentanil, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus or tacrolimus).

Imatinib mesylate will increase plasma concentration of other CYP3A4 metabolized drugs (e.g., triazolo-benzodiazepines, dihydropyridine calcium channel blockers, certain HMG-CoA reductase inhibitors, etc.).

Because warfarin is metabolized by CYP2C9 and CYP3A4, patients who require anticoagulation should receive low-molecular weight or standard heparin instead of warfarin.

Interactions with Drugs Metabolized by CYP2D6

Imatinib mesylate increases the mean C_{max} and AUC of metoprolol by approximately 23% suggesting that Imatinib mesylate has a weak inhibitory effect on CYP2D6-mediated metabolism. No dose adjustment is necessary, however, caution is recommended when administering Imatinib mesylate with CYP2D6 substrates that have a narrow therapeutic window.

Interactions with Acetaminophen

In vitro, Imatinib mesylate inhibits the acetaminophen O-glucuronidate pathway (*K*i 58.5 µM). Coadministration of Imatinib mesylate (400 mg/day for 8 days) with acetaminophen (1000 mg single dose on day 8) in patients with CML did not result in any changes in the pharmacokinetics of acetaminophen. Imatinib mesylate plasma protein binding was not altered in the presence of single-dose acetaminophen. There is no pharmacokinetic or safety data on the concomitant use of Imatinib mesylate at doses greater than 400 mg/day or the chronic use of concomitant acetaminophen and Imatinib mesylate.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In the 2-year rat carcinogenicity study administration of imatinib at 15, 30, and 60 mg/kg/day resulted in a statistically significant reduction in the longevity of males at 60 mg/kg/day and females at greater than or equal to 30 mg/kg/day. Target organs for neoplastic changes were the kidneys (renal tubule and renal pelvis), urinary bladder, urethra, preputial and clitoral gland, small intestine, parathyroid glands, adrenal glands and non-glandular stomach. Neoplastic lesions were not seen at 30 mg/kg/day for the kidneys, urinary bladder, urethra, small intestine, parathyroid glands, adrenal glands and non-glandular stomach, and 15 mg/kg/day for the preputial and clitoral gland. The papilloma/carcinoma of the preputial/clitoral gland were noted at 30 and 60 mg/kg/day, representing approximately 0.5 to 4 or 0.3 to 2.4 times the human exposure (based on AUC) at 400 mg/day or 800 mg/day, respectively, and at 0.4 to 3.0 times the drug exposure in children (based on AUC) at 340 mg/m². The renal tubule adenoma/carcinoma, renal pelvis transitional cell neoplasms, the urinary bladder and urethra transitional cell papillomas, the small intestine adenocarcinomas, the parathyroid glands adenomas, the benign and malignant medullary tumors of the adrenal glands and the non-glandular stomach papillomas/carcinomas were noted at 60 mg/kg/day. The relevance of these findings in the 2-year carcinogenicity study for humans is not known. Positive genotoxic effects were obtained for imatinib in an *in vitro* mammalian cell assay (Chinese hamster ovary cells, clastogenicity (chromosome aberrations)) in the presence and absence of metabolic activation. Two intermediates of the manufacturing process, which are also present in the final product, are positive for mutagenesis in the Ames assay. One of these intermediates was also positive in the mouse lymphoma assay. Imatinib was not genotoxic when tested in an *in vitro* bacterial cell assay (Ames test), an *in vitro* mammalian cell assay (mouse lymphoma) and an *in vivo* micronucleus assay.

In a study of fertility, male rats were dosed for 70 days prior to mating and female rats were dosed 14 days prior to mating and through to gestational Day 6. Testicular and epididymal weights and percent motile sperm were decreased at 60 mg/kg, approximately one-half the maximum human dose based on body surface area. Fertility of males or females was not seen at doses less than or equal to 20 mg/kg (one-fourth the maximum human dose of 800 mg). The fertility of male and female rats not affected.

Fertility was not affected in the preclinical fertility and early embryonic development study although lower testes and epididymis weights, the urinary bladder and urethra transitional cell papillomas, the small intestine adenocarcinomas, the parathyroid glands adenomas, the benign and malignant medullary tumors of the adrenal glands and the non-glandular stomach papillomas/carcinomas were noted at 60 mg/kg/day. The relevance of these findings in the 2-year carcinogenicity study for humans is not known. Positive genotoxic effects were obtained for imatinib in an *in vitro* mammalian cell assay (Chinese hamster ovary cells, clastogenicity (chromosome aberrations)) in the presence and absence of metabolic activation. Two intermediates of the manufacturing process, which are also present in the final product, are positive for mutagenesis in the Ames assay. One of these intermediates was also positive in the mouse lymphoma assay. Imatinib was not genotoxic when tested in an *in vitro* bacterial cell assay (Ames test), an *in vitro* mammalian cell assay (mouse lymphoma) and an *in vivo* micronucleus assay.

13.2 Animal Toxicology and/or Pharmacology

Toxicities from Long-Term Use

It is important to consider potential toxicities suggested by animal studies, specifically, *liver, kidney, and cardiac toxicity and immunosuppression*. Severe liver toxicity was observed in dogs treated for 2 weeks, with elevated liver enzymes, hepatocellular necrosis, bile duct necrosis, and bile duct hyperplasia. Renal toxicity was observed in monkeys treated for 2 weeks, with focal mineralization and dilation of the renal tubules and tubular nephrosis. Increased BUN and creatinine were observed in several of these animals. An increased rate of opportunistic infections was observed with chronic imatinib treatment in laboratory animal studies. In a 39 week monkey study, treatment with imatinib resulted in worsening of normally diagnosed tuberculosis infections in these animals. Lymphoma was observed in animals (as in humans). Additional long-term toxicities were identified in a 2-year rat study. Histopathological examination of the rats 28 months after dosing with imatinib *(cardiomyopathy* (both cases), chronic progressive nephropathy (females) and the preputial gland papilloma as principal causes of death or reasons for sacrifice. Non-neoplastic lesions seen in this 2-year study which were not identified in earlier preclinical studies were cardiovascular system; pancreas, exocrine glands and teeth. The most important changes included cardiac hypertrophy and dilatation, leading to signs of cardiac insufficiency in some animals.

14 CLINICAL STUDIES

14.1 Chronic Myeloid Leukemia

Chronic Phase, Newly Diagnosed:

An open-label, multicenter, international randomized Phase 3 study (Imatinib mesylate versus INF-α+Ara-C) has been conducted in patients with newly diagnosed Philadelphia chromosome positive (Ph+) chronic myeloid leukemia (CML) in chronic phase. This study compared treatment with either single-agent Imatinib mesylate or a combination of interferon-alpha (IFN) plus cytarabine (Ara-C). Patients were allowed to cross over to the alternative treatment arm if they failed to show a complete hematologic response (CHR) at 6 months, a major cytogenetic response (MCyR) at 12 months, or if they lost a CHR or MCyR. Patients with increasing white blood cell counts (leukemic progression) were also allowed to cross over to the alternative treatment arm with the permission of the study monitoring committee (SMC). In the Imatinib mesylate arm, patients were initially treated with 400 mg daily. Dose escalations were allowed from 400 mg daily to 600 mg daily, then from 600 mg daily to 800 mg daily. In the IFN arm, patients were treated with a target dose of IFN of 5 MIU/m² daily subcutaneously in combination with subcutaneous Ara-C 20 mg/m²/day for 10 days/month.

A total of 1,106 patients were randomized from 177 centers in 16 countries, 553 to each arm. Baseline characteristics were well balanced between the two arms. Median age was 51 years (range 18 to 70 years), with 21.9% of patients greater than or equal to 65 years old. There were 59% males and 41% females. 89.9% Caucasian and 4.1% Black patients. At the cut-off for this analysis (7 years after last patient had been recruited), the median duration of first-line treatment was 82 and 8 months in the Imatinib mesylate and IFN arm, respectively. The median duration of second-line treatment with Imatinib mesylate was 64 months. Sixty percent of patients randomized to Imatinib mesylate are still receiving first-line treatment. In these patients, the average dose of Imatinib mesylate was 419 mg/day ± 57 mg. Overall, in patients receiving first line Imatinib mesylate, the average daily dose delivered was 406 mg ± 76 mg. Due to discontinuations and crossovers, only 2% of patients randomized to IFN were still on first-line treatment. In the IFN arm, withdrawal of consent (14%) was the most frequent reason for discontinuation of first-line therapy, and the most frequent reason for cross over to the Imatinib mesylate arm was severe intolerance to treatment (26%) and progression (14%).

The primary efficacy endpoint of the study was progression-free survival (PFS). Progression was defined as any of the following events: progression to accelerated phase or blast crisis (AP/BC), death, loss of CHR or MCyR, or in patients not achieving a CHR an increasing WBC despite appropriate therapeutic management. The protocol specified that the progression analysis would compare the intent to treat (ITT) Population: patients randomized to receive Imatinib mesylate were compared with patients randomized to receive IFN. Patients that crossed over prior to progression were censored at the time of cross-over, and events that occurred in these patients following cross-over were attributed to the original randomized treatment. The estimated rate of progression-free survival at 84 months in the ITT population was 81.2 % (95% CI: 78, 83) in the Imatinib mesylate arm and 60.6 % (56, 65) in the IFN arm (p less than 0.0001, log-rank test). (Figure 1). With 7 years follow up there were 93 (16.8%) progression events in the Imatinib mesylate arm: 37(6.7%) progression to AP/BC, 31(5.6%) loss of MCyR, 15 (2.7%) loss of CHR or increase in WBC and 10 (1.8%) CML unrelated deaths. In contrast, there were 165 (29.8%) events in the IFN-Ara-C arm of which 130 occurred during first-line treatment. In the IFN arm, the estimated rate of patients free of progression to accelerated phase (AP) or blast crisis (BC) at 84 months was 92.5%(96, 95) in the Imatinib mesylate arm compared to the 85.1%, (82, 89) (p less than or equal to 0.001) in the IFN arm, (Figure 2). The annual rates of any progression events have decreased with time on therapy. The probability of progression free at 84 months was 85% for patients who were in complete cytogenetic response (CCyR) with molecular response (greater than or equal to 3 log reduction in BCR-ABL transcripts as measured by quantitative reverse transcriptase polymerase chain reaction) at 12 months, compared to 89% for patients in complete cytogenetic response but without a major molecular response and 70% in patients who were not in complete cytogenetic response at this time point (p less than 0.001).

Figure 1: Progression Free Survival (ITT Principle)



Figure 2: Time to Progression to AP or BC (ITT Principle)



A total of 71 (12.8%) and 85 (15.4%) patients died in the Imatinib mesylate and IFN+Ara-C group, respectively. At 84 months the estimated survival rate was 86.4% (83, 90) in the Imatinib (400, 87) in the randomized Imatinib mesylate and IFN+Ara-C group, respectively (p=0.073 log-rank test). The hazard ratio is 0.750 with 95% CI 0.547 to 1.028. This time-to-event endpoint may be affected by the high crossover rate from IFN+Ara-C to Imatinib mesylate. Major cytogenetic response, hematologic response, evaluation of minimal residual disease (molecular response), time to accelerated phase or blast crisis and survival were main secondary endpoints. Response data are shown in Table 18. Complete hematologic response, major cytogenetic response and complete cytogenetic response were also statistically significantly higher in patients receiving Imatinib mesylate at 84 months vs. IFN+Ara-C arm (no cross-over data considered for evaluation of response). Median time to CCyR in the 454 responders was 6 months (range 2 to 64 months, 25th to 75th percentiles=3 to 11 months). 90% of responders seen after 22 months of therapy.

Table 18: Response in Newly Diagnosed CML Study (84-Month Data)

(Best Response Rate)	Imatinib mesylate n=553	IFN+Ara-C n=553
Hematologic Response[a]		
CHR Rate n (%)	534 (96.6%)* (94.7%, 97.9%)	313 (56.6%)* (52.4%, 60.8%)
CCyR n (%)	472 (85.4 %)* (82.1%, 88.2%)	93 (16.8%)* (13.8%, 20.2%)
Cytogenetic Response[b]		
Major Cytogenetic Response n (%)	473 (74.7%)* (70.8, 78.3)	36 (6.5%)* (4.6, 8.9)
Unconfirmed[c]	82.5%*	11.6%*

[a] less than 0.001, Fischer's exact test

Hematologic response criteria (all responses to be confirmed after greater than or equal to 4 weeks): WBC less than 10 x 10⁹/L, platelet less than 450 x 10⁹/L, myelocyte + metamyelocyte less than 5% in blood, no basilic/promyelocytes in blood, basophils less than 20%, no extramedullary involvement).

Cytogenetic response criteria (confirmed after greater than or equal to 4 weeks): complete (0% Ph+ metaphases) or partial (1% to 35%). A major response (0% to 35% combines both complete and partial responses.

Unconfirmed cytogenetic response is based on a single bone marrow cytogenetic evaluation, therefore unconfirmed complete or partial cytogenetic responses might have had a lesser cytogenetic response on a subsequent bone marrow evaluation.

Molecular response was defined as follows: in the peripheral blood, after 12 months of therapy, reduction of greater than or equal to 3 logarithms in the amount of BCR-ABL transcripts (measured by real-time quantitative reverse transcriptase polymerase chain reaction) normalized to standardized baseline. Molecular response was only evaluated in a subset of patients who had a complete cytogenetic response by 12 months or later (N=333). The molecular response rate in patients who had a complete cytogenetic response in the Imatinib mesylate arm was 59% at 12 months and 72% at 24 months.

Physical, functional, and treatment-specific biologic response modifier scales from the FACT-BMT (Functional Assessment of Cancer Therapy - Biologic Response Modifier) instrument were used to assess patient-reported general well-being in patients treated at CML clinics. Patients with CML in chronic phase. After one month of therapy to 6 months of therapy, there was a 13% to 21% decrease in median index from baseline in patients treated with IFN, consistent with increased symptoms of IFN toxicity. There was an improvement change from baseline in median index for patients treated with Imatinib mesylate.

An open-label, multicenter, randomized trial (Imatinib mesylate versus nilotinib) was conducted to determine the efficacy of Imatinib mesylate versus nilotinib in adult patients with cytogenetically confirmed, newly diagnosed Ph+ CML-CP. Patients within 6 months of diagnosis and were previously untreated for CML-CP, except for hydroxyurea and/or anagrelide. Efficacy was based on a total of 846 patients; 283 patients in the Imatinib mesylate 400 mg once daily group, 282 patients in the nilotinib 300 mg twice daily group, and 281 patients in the nilotinib 400 mg twice daily group.

Median age was 46 years in the Imatinib mesylate group and 47 years in both nilotinib groups, with 12%, 13%, and 10% patients greater than or equal to 65 years of age in Imatinib mesylate 400 mg once-daily, nilotinib 300 mg twice daily and nilotinib 400 mg twice daily treatment groups, respectively. There were slightly more male than female patients in all groups (56%, 56%, and 62% in Imatinib mesylate 400 mg once-daily, nilotinib 300 mg twice-daily and nilotinib 400 mg twice-daily treatment groups, respectively). More than 60% of patients were Caucasian, and 22% were Asian.

The primary data analysis was performed when all 846 patients completed 12 months of treatment or discontinued earlier. Subsequent analyses were done when patients completed 24, 36 and 48 months of treatment or discontinued earlier. The median time on treatment was approximately 61 months in all three treatment groups.

The primary efficacy endpoint was major molecular response (MMR) at 12 months after the start of study medication. MMR was defined as less than or equal to 0.1% BCR-ABL/ABL % by international scale measured by RQ-PCR, which corresponds to a greater than or equal to 3 log reduction of BCR-ABL transcript from standardized baseline. Efficacy endpoints are summarized in Table 19.

Twelve patients in the Imatinib mesylate arm progressed to either accelerated phase or blast crises (7 patients within first 6 months, 2 patients within 6 to 12 months, 2 patients within 12 to 18 months and 1 patient within 18 to 24 months) while two patients on the nilotinib arm progressed to either accelerated phase or blast crisis (both within the first 6 months of treatment).

Table 19: Efficacy (MMR and CCyR) at Imatinib Mesylate Compared to Nilotinib in Newly Diagnosed Ph+ CML-CP

	Imatinib mesylate 400 mg once daily	nilotinib 300 mg twice daily
	N=283	N=282
MMR at 12 months (95% CI)	22% (17.6, 27.6)	44% (38.4, 50.3)
P-Value[a]		< 0.0001
CCyR by 12 months (95% CI)	65% (59.2, 70.6)	80% (75.0, 84.6)
MMR at 24 months (95% CI)	38% (31.8, 43.4)	62% (55.8, 67.4)
CCyR[b] by 24 months (95% CI)	77% (71.7, 81.8)	87% (82.4, 90.6)

[a]CMH test stratified by Sokal risk group

[b]CCyR: 0% Ph+ metaphases. Cytogenetic responses were based on the percentage of Ph-positive metaphases among greater than or equal to 20 metaphase cells in each bone marrow sample.

By the 60 months, MMR was achieved by 60% of patients on Imatinib mesylate and 77% of patients on nilotinib.

Median overall survival was not reached in either arm. At the time of the 60-month final analysis, the estimated survival rate was 91.7% for patients on Imatinib mesylate and 93.7% for patients on nilotinib.

Late Chronic Phase CML and Advanced Stage CML: Three international, open-label, single-arm Phase 2 studies were conducted to determine the safety and efficacy of Imatinib mesylate in patients with Ph+ CML: 1) in the chronic phase after failure of IFN therapy, 2) in accelerated phase disease, or 3) in myeloid blast crisis. About 42% of patients were women and 8% were black. In clinical studies, 38% to 40% of patients were greater than or equal to 65 years of age and 10% to 12% of patients were greater than or equal to 70 years of age.

Chronic Phase, Prior Interferon-Alpha Treatment: 532 patients were treated at a starting dose of 400 mg; dose escalation to 600 mg was allowed. The patients were distributed in three main categories according to their response to prior interferon: failure