

<b>Concord Biotech Limited</b> 297-298/2P, Valthera, Tal. Dholka, Dist. Ahmedabad - 382 225							
<b>Artwork Approval: Medication Guide</b>							
<b>Product:</b>	Mycophenolate Mofetil Capsules, USP 250 mg	<b>Market:</b>	US	<b>Size:</b>	220 x 470 mm	<b>Artwork Supersede:</b>	VP00209-01
<b>Item code:</b>	VP00209-02	<b>Mfg. Location:</b>	Valthera	<b>Com. &amp; Vendor:</b>	Medication Guide / Temple	<b>P.code:</b>	NA
<b>Substrate:</b>	40 gsm bible paper printed Medication guide for Mycophenolate mofetil capsule USP 250mg						
<b>Reason for Issuance:</b>	Artworks revised for toll free number						

**220.00 mm**

470.00 mm

**MEDICATION GUIDE**

**Mycophenolate Mofetil Capsules, USP 250 mg  
(MYE koe FEN oh late MOE fe til)**

Read the Medication Guide that comes with Mycophenolate mofetil capsules before you start taking it and each time you refill your prescription. There may be new information. This Medication Guide does not take the place of talking with your doctor about your medical condition or your treatment.

**What is the most important information I should know about mycophenolate mofetil capsules?**  
**Mycophenolate mofetil capsules can cause serious side effects, including:**  
**Increased risk of loss of a pregnancy (miscarriage) and higher risk of birth defects.** Females who take mycophenolate mofetil during pregnancy have a higher risk of miscarriage during the first 3 months (first trimester), and a higher risk that their baby will be born with birth defects.

- If you are a female who can become pregnant,** your doctor must talk with you about acceptable birth control methods (contraceptive counseling) to use while taking mycophenolate mofetil capsules. You should have 1 pregnancy test immediately before starting mycophenolate mofetil capsules and another pregnancy test 8 to 10 days later. Pregnancy tests should be repeated during routine follow-up visits with your doctor. Talk to your doctor about the results of all of your pregnancy tests. You must use acceptable birth control during your entire mycophenolate mofetil treatment and for 6 weeks after stopping mycophenolate mofetil, unless at any time you choose to avoid sexual intercourse (abstinence) with a man completely. Mycophenolate mofetil decreases blood levels of the hormones in birth control pills that you take by mouth. Birth control pills may not work as well while you take mycophenolate mofetil capsules, and you could become pregnant. If you take birth control pills while using mycophenolate mofetil capsules you must also use another form of birth control. Talk to your doctor about other birth control methods that you can use while taking mycophenolate mofetil capsules.
- If you are sexually active male whose female partner can become pregnant while you are taking mycophenolate mofetil capsules,** use effective contraception during treatment and for at least 90 days after stopping mycophenolate mofetil capsules.
- If you plan to become pregnant,** talk with your doctor. Your doctor will decide if other medicines to prevent rejection may be right for you.
- If you become pregnant while taking mycophenolate mofetil capsules, do not stop taking mycophenolate mofetil capsules. Call your doctor right away.** You and your doctor may decide that other medicines to prevent rejection may be right for you. You and your doctor should report your pregnancy to the Mycophenolate Pregnancy Registry either:
  - By phone at 1-800-617-8191 or
  - By visiting the REMS website at [www.mycophenolateREMS.com](http://www.mycophenolateREMS.com)

The purpose of this registry is to gather information about the health of you and your baby.

**Increased risk of getting certain cancers.** People who take Mycophenolate mofetil have a higher risk of getting lymphoma, and other cancers, especially skin cancer. Tell your doctor if you have:

- unexplained fever, prolonged tiredness, weight loss or lymph node swelling
- a brown or black skin lesion with uneven borders, or one part of the lesion does not look like the other
- a change in the size and color of a mole
- a new skin lesion or bump
- any other changes to your health

**Increased risk of getting serious infections.** Mycophenolate mofetil weakens the body's immune system and affects your ability to fight infections. Serious infections can happen with mycophenolate mofetil capsules and can lead to hospitalizations and death. These serious infections can include:

- Viral infections.** Certain viruses can live in your body and cause active infections when your immune system is weak. **Viral infections that can happen with mycophenolate mofetil include:**
  - Shingles, other herpes infections, and cytomegalovirus (CMV). CMV can cause serious tissue and blood infections.
  - BK virus. BK virus can affect how your kidney works and cause your transplanted kidney to fail.
  - Hepatitis B and C viruses. Hepatitis viruses can affect how your liver works. Talk to your doctor about how hepatitis viruses may affect you.
- A brain infection called Progressive Multifocal Leukoencephalopathy (PML).** In some patients,

mycophenolate mofetil may cause an infection of the brain that may cause death. You are at risk for this brain infection because you have a weakened immune system. Call your doctor right away if you have any of the following symptoms:

- Weakness on one side of the body
- You do not care about things that you usually care about (apathy)
- You are confused or have problems thinking
- You can not control your muscles

- Fungal infections.** Yeasts and other types of fungal infections can happen with mycophenolate mofetil and can cause serious tissue and blood infections (see "**What are the possible side effects of mycophenolate mofetil capsules?**").

**Call your doctor right away if you have any of the following signs and symptoms of infection:**

- temperature of 100.5°F or greater
- cold symptoms, such as a runny nose or sore throat
- flu symptoms, such as an upset stomach, stomach pain, vomiting or diarrhea
- earache or headache
- pain during urination
- white patches in the mouth or throat
- unexpected bruising or bleeding
- cuts, scrapes or incisions that are red, warm and oozing pus

See "**What are the possible side effects of mycophenolate mofetil capsules?**" for information about other serious side effects.

**What is mycophenolate mofetil capsules?**

- Mycophenolate mofetil capsules is a prescription medicine to prevent rejection (antirejection medicine) in people who have received a kidney, heart or liver transplant. Rejection is when the body's immune system perceives the new organ as a "foreign" threat and attacks it.
- Mycophenolate mofetil capsules is used with other medicines containing cyclosporine and corticosteroids.

**Who should not take mycophenolate mofetil capsules?**  
**Do not take mycophenolate mofetil capsules if you are allergic to mycophenolate mofetil or any of the ingredients in mycophenolate mofetil capsules.** See the end of this Medication Guide for a complete list of ingredients in mycophenolate mofetil capsules.

**What should I tell my doctor before taking mycophenolate mofetil capsules?**  
**Tell your doctor about all of your medical conditions, including if you:**

- have any digestive problems, such as ulcers.
- have Lesch-Nyhan or Kelley-Seegmiller syndrome or another rare inherited deficiency hypoxanthine-guanine phosphoribosyl-transferase (HGPRT). You should not take mycophenolate mofetil capsules if you have one of these disorders.
- plan to receive any vaccines. People taking mycophenolate mofetil capsules should not receive live vaccines. Some vaccines may not work as well during treatment with mycophenolate mofetil capsules.
- are pregnant or plan to become pregnant. See "**What is the most important information I should know about mycophenolate mofetil capsules?**"
- are breastfeeding or plan to breastfeed. It is not known if mycophenolate mofetil passes into breast milk. You and your doctor will decide if you will take mycophenolate mofetil capsules or breastfeed.

**Tell your healthcare provider about all of the medicines you take,** including prescription and over-the-counter medicines, vitamins and herbal supplements. Some medicines may affect the way mycophenolate mofetil works, and mycophenolate mofetil may affect how some medicines work. Especially tell your doctor if you take:

- birth control pills (oral contraceptives). See "**What is the most important information I should know about mycophenolate mofetil capsules?**"
- sevelamer (Renagel<sup>®</sup>, Renvela<sup>™</sup>). These products should be taken at least 2 hours after taking mycophenolate mofetil capsules
- acyclovir (Zovirax<sup>®</sup>), valacyclovir (Valtrex<sup>®</sup>), ganciclovir (CYTOVENE<sup>®</sup>-IV, Vitrasert<sup>®</sup>), valganciclovir (VALCYTE<sup>®</sup>)
- rifampin (Rifater<sup>®</sup>, Rifamate<sup>®</sup>, Rimactane<sup>®</sup>, Rifadin<sup>®</sup>)
- antacids that contain magnesium and aluminum (mycophenolate mofetil capsules and the antacid should not be taken at the same time)
- proton pump inhibitors (PPIs) (Prevacid<sup>®</sup>, Protonix<sup>®</sup>)
- sulfamethoxazole/trimethoprim (BACTRIM<sup>™</sup>, BACTRIM Ds<sup>™</sup>)
- norfloxacin (Noroxin<sup>®</sup>) and metronidazole (Flagyl<sup>®</sup>, Flagyl<sup>®</sup> ER, Flagyl<sup>®</sup> IV, Metro IV, Helidac<sup>®</sup>, Pylera<sup>™</sup>)
- ciprofloxacin (Cipro<sup>®</sup>, Cipro<sup>®</sup> XR, Ciloxan<sup>®</sup>, Proquin<sup>®</sup> XR) and amoxicillin plus clavulanic acid (Augmentin<sup>®</sup>,

Front

Front

Folded Size : 220 x 240 mm

**PAN Process Black C**

**MG MMC US LUPIN\_02\_05\_2019\_01**

<b>Prepared By</b>	<b>Reviewed By</b>	<b>Reviewed By</b>	<b>Reviewed By</b>	<b>Reviewed By</b>	<b>Approved By</b>
<b>Pack F&amp;D</b>	<b>F&amp;D</b>	<b>RA</b>	<b>Production</b>	<b>QC</b>	<b>QA</b>



**Artwork Approval: Medication Guide**

<b>Product:</b>	Mycophenolate Mofetil Capsules, USP 250 mg	<b>Market:</b>	US	<b>Size:</b>	220 x 470 mm	<b>Artwork Supersede:</b>	VP00209-01	
<b>Item code:</b>	VP00209-02	<b>Mfg. Location:</b>	Valthera	<b>Com. &amp; Vendor:</b>		Medication Guide / Temple	<b>P.code:</b>	NA
<b>Substrate:</b>	40 gsm bible paper printed Medication guide for Mycophenolate mofetil capsule USP 250mg							
<b>Reason for Issuance:</b>	Artworks revised for toll free number							

**220.00 mm**

**470.00 mm**

<p>Augmentin Xr™)</p> <ul style="list-style-type: none"> <li>• azathioprine (Azasan®, Imuran®)</li> <li>• cholestyramine (Questran Light®, Questran®, Locholest Light, Locholest, Prevalite®)</li> </ul> <p>Know the medicines you take. Keep a list of them to show to your doctor or nurse and pharmacist when you get a new medicine. Do not take any new medicine without talking with your doctor.</p> <p><b>How should I take mycophenolate mofetil capsules?</b></p> <ul style="list-style-type: none"> <li>• Take mycophenolate mofetil capsules exactly as prescribed.</li> <li>• Do not stop taking mycophenolate mofetil capsules or change the dose unless your doctor tells you to.</li> <li>• If you miss a dose of mycophenolate mofetil capsules, or you are not sure when you took your last dose, take your prescribed dose of mycophenolate mofetil capsules as soon as you remember. If your next dose is less than 2 hours away, skip the missed dose and take your next dose at your normal scheduled time. Do not take 2 doses at the same time. Call your doctor if you are not sure what to do.</li> <li>• Take mycophenolate mofetil capsules on an empty stomach, unless your doctor tells you otherwise.</li> <li>• <b>Do not</b> open or crush mycophenolate mofetil capsules.</li> <li>• If you are not able to swallow mycophenolate mofetil capsules, your doctor may prescribe mycophenolate mofetil Oral Suspension. This is a liquid form of mycophenolate mofetil. Your pharmacist will mix the medicine before you pick it up from a pharmacy.</li> <li>• <b>Do not</b> breath in (inhale) or let mycophenolate mofetil powder come in contact with your skin or mucous membranes.             <ul style="list-style-type: none"> <li>• If you accidentally get the powder on the skin, wash the area well with soap and water.</li> <li>• If you accidentally get the powder in your eyes or other mucous membranes, flush with plain water.</li> </ul> </li> <li>• If you take too much mycophenolate mofetil capsules, call your doctor or the poison control center right away.</li> </ul> <p><b>What should I avoid while taking mycophenolate mofetil capsules?</b></p> <ul style="list-style-type: none"> <li>• Avoid becoming pregnant. See "<b>What is the most important information I should know about mycophenolate mofetil capsules?</b>"</li> <li>• Limit the amount of time you spend in sunlight. Avoid using tanning beds or sunlamps. People who take mycophenolate mofetil capsules have a higher risk of getting skin cancer. (See "<b>What is the most important information I should know about mycophenolate mofetil capsules?</b>") Wear protective clothing when you are in the sun and use a sunscreen with a high protection factor. This is especially important if your skin is very fair or if you have a family history of skin cancer.</li> <li>• You should not donate blood while taking mycophenolate mofetil capsules and for at least 6 weeks after stopping mycophenolate mofetil capsules.</li> <li>• You should not donate sperm while taking mycophenolate mofetil capsules and for 90 days after stopping mycophenolate mofetil capsules.</li> </ul> <ul style="list-style-type: none"> <li>• Mycophenolate mofetil capsules may influence your ability to drive and use machines (See "<b>What are the possible side effects of mycophenolate mofetil capsules?</b>"). If you experience drowsiness, confusion, dizziness, tremor, or low blood pressure during treatment with mycophenolate mofetil capsules, you should be cautious about driving or using heavy machines.</li> </ul> <p><b>What are the possible side effects of mycophenolate mofetil capsules?</b> <b>Mycophenolate mofetil capsules can cause serious side effects, including:</b></p> <ul style="list-style-type: none"> <li>• See "<b>What is the most important information I should know about mycophenolate mofetil capsules?</b>"</li> <li>• <b>Low blood cell counts.</b> People taking high doses of mycophenolate mofetil capsules each day may have a decrease in blood counts, including:             <ul style="list-style-type: none"> <li>• <b>white blood cells, especially neutrophils.</b> Neutrophils fight against bacterial infections. You have a higher chance of getting an infection when your white blood cell count is low. This is most common from 1 month to 6 months after your transplant.</li> <li>• <b>red blood cells.</b> Red blood cells carry oxygen to your body tissues. You have a higher chance of getting severe anemia when your red blood cell count is low.</li> <li>• <b>platelets.</b> Platelets help with blood clotting. Your doctor will do blood tests before you start taking mycophenolate mofetil capsules and during treatment with mycophenolate mofetil capsules to check your blood cell counts. Tell your doctor right away if you have any signs of infection (see "<b>What is the most important information I should know about mycophenolate mofetil capsules?</b>"), including any unexpected bruising or bleeding. Also, tell your doctor if you have unusual tiredness, lack of energy, dizziness or fainting.</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• <b>Stomach problems.</b> Stomach problems including intestinal bleeding, a tear in your intestinal wall (perforation) or stomach ulcers can happen in people who take mycophenolate mofetil capsules. Bleeding can be severe and you may have to be hospitalized for treatment. Call your doctor right away if you have sudden or severe stomach-area pain or stomach-area pain that does not go away, or if you have diarrhea.</li> </ul> <p><b>The most common side effects of mycophenolate mofetil capsules include:</b></p> <ul style="list-style-type: none"> <li>• diarrhea</li> <li>• blood problems including low white and red blood cell counts</li> <li>• infections</li> <li>• blood pressure problems</li> <li>• fast heart beat</li> <li>• swelling of the lower legs, ankles and feet</li> <li>• changes in laboratory blood levels, including high levels of blood sugar (hyperglycemia)</li> <li>• stomach problems including diarrhea, constipation, nausea and vomiting</li> <li>• rash</li> <li>• nervous system problems such as headache, dizziness and tremor</li> </ul> <p><b>Side effects that happen more often in children than in adults taking mycophenolate mofetil capsules include:</b></p> <ul style="list-style-type: none"> <li>• stomach area pain</li> <li>• fever</li> <li>• infection</li> <li>• pain</li> <li>• blood infection (sepsis)</li> <li>• diarrhea</li> <li>• vomiting</li> <li>• sore throat</li> <li>• colds (respiratory tract infections)</li> <li>• high blood pressure</li> <li>• low white blood cell count</li> <li>• low red blood cell count</li> </ul> <p>These are not all of the possible side effects of mycophenolate mofetil capsules. Tell your doctor about any side effect that bothers you or that does not go away. <b>Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.</b></p> <p><b>You may also report side effects to Concord Biotech Limited at Telephone : 1-844-553-5534, Fax : 1-844-552-5515.</b></p> <p><b>How should I store mycophenolate mofetil capsules?</b></p> <ul style="list-style-type: none"> <li>• Store mycophenolate mofetil capsules at room temperature, between 59°F to 86°F (15°C to 30°C).</li> </ul> <p><b>Keep mycophenolate mofetil capsules and all medicines out of the reach of children.</b></p> <p><b>General Information about the safe and effective use of mycophenolate mofetil capsules.</b> Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use mycophenolate mofetil capsules for a condition for which it was not prescribed. Do not give mycophenolate mofetil capsules to other people, even if they have the same symptoms that you have. It may harm them. This Medication Guide summarizes the most important information about mycophenolate mofetil capsules. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about mycophenolate mofetil capsules that is written for health professionals.</p> <p><b>What are the ingredients in mycophenolate mofetil capsules?</b> <b>Active Ingredient:</b> mycophenolate mofetil <b>Inactive Ingredients:</b> microcrystalline cellulose, hydroxypropyl cellulose, povidone (K-90), croscarmellose sodium, talc and magnesium stearate. The capsule shells contain FD&amp;C blue #2, gelatin, red iron oxide, sodium lauryl sulfate, titanium dioxide and yellow iron oxide. The imprinting ink contains shellac, dehydrated alcohol, isopropyl alcohol, butyl alcohol, propylene glycol, strong ammonia solution, iron oxide black and potassium hydroxide.</p> <p>Manufactured By: <b>Concord Biotech Limited,</b> 297-298/2P, At Valthera Ta. Dholka, Ahmedabad – 382225, Gujarat, India.</p> <p>Manufactured for: <b>Lupin Pharmaceuticals, Inc.</b> Baltimore, Maryland 21202 United States</p> <p>Any other trademarks in this document are the property of their respective owners.</p> <p>This Medication Guide has been approved by the U.S. Food and Drug Administration.</p> <p style="text-align: right;">MG Revised: May 2019</p> <p>VP00209-02</p>
--	---

Back

Folded Size : 220 x 240 mm

Back

**PAN Process Black C**

**MG\_MMC\_US\_LUPIN\_02\_05\_2019\_01**

Prepared By	Reviewed By	Reviewed By	Reviewed By	Reviewed By	Approved By
Pack F&D	F&D	RA	Production	QC	QA