

WARNING: ADDICTION, ARUSE, AND MISUSE: RISK EVALUATION AND MITIGATION STRATEGY (REMS): LIFE-THREATENING RESPIRATORY DEPRESSION: ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

<u>Addiction, Abuse, and Misuse</u>
Pentazocine and Naloxone Tablets exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing Pentazocine and Naloxone Tablets, and monitor all patients regularly for the development of these behaviors and conditions [see WARNINGS]

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS):
To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse,
the Food and Drug Administration (FDA) has required a REMS for these products [see Warnings]. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare provider

- are strongly encouraged to complete a REMS-compliant education program, counsel patients and/or their caregivers, with every prescription, on safe use, serious risks, storage, and disposal of these products.
- emphasize to patients and their caregivers the importance of reading the Medication Guide every time it is provided by their pharmacist, and consider other tools to improve patient, household, and community safety.

### Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of Pentazocine and Naloxone Tablets. Monitor for respiratory depression, especially during initiation of Pentazocine and Naloxone Tablets or following a dose increase [see WARNINGS].

### Accidental Ingestion

Accidental innestion of even one dose of Pentazocine and Naloxone Tablets, especially by children can result in a fatal overdose of Pentazocine [see WARNINGS].

Neonatal Opioid Withdrawal Syndrome Prolonged use of Pentazocine and Naloxone Tablets during pregnancy can result in neonatal opioic withdrawal syndrome, which may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see WARNINGS].

## Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants

ant use of opioids with benzodiazenines or other central nervous system (CNS) depressant including alcohol, may result in profound sedation, respiratory depression, coma, and death [see Warnings and Precautions, Drug Interactions].


Reserve concomitant prescribing of Pentazocine and Naloxone Tablets and benzodiazepines or

- other CNS depressants for use in patients for whom alternative treatment options are inadequate Limit dosages and durations to the minimum required.
- Follow patients for signs and symptoms of respiratory depression and sedation.


Pentazocine and Naloxone Tablets, USP contain pentazocine hydrochloride, USP, a partial opioid agonist equivalent to 50 mg base and is a member of the benzazocine series (also known as the benzom series), and naloxone hydrochloride, USP, an opioid antagonist equivalent to 0.5 mg base.

Pentazocine and Naloxone Tablets, USP are an analgesic for oral administration.

Chemically, pentazocine hydrochloride, USP is  $(2R^*,6R^*,11R^*)$ -1,2,3,4,5,6-Hexahydro-6,11-dimethyl-3-(3-methyl-2-butenyl)-2,6-methano-3-benzazocin-8-ol hydrochloride, a white, crystalline substance soluble in acidic aqueous solutions, and has the following structural formula:



Chemically, naloxone hydrochloride, USP is Morphinan-6-one,4,5-epoxy-3,14-dihydroxy-17-(2propenyl)-hydrochloride,  $(5\alpha)$ -. It is a slightly off-white powder, and is soluble in water and dilute acids, and has the following structural formula:



Inactive Ingredients: colloidal silicon dioxide, dibasic calcium phosphate, D&C Yellow No. 10, magnesium stearate, microcrystalline cellulose, sodium lauryl sulfate and pregelatinized starch

# CLINICAL PHARMACOLOGY

Pentazocine is a mixed agonist-antagonist at opioid receptors. Pentazocine is a partial agonist at the mu

Naloxone is an opioid antagonist.

# **Effects on the Central Nervous System**

Pentazocine produces respiratory depression by direct action on brain stem respiratory centers. The respiratory depression involves a reduction in the responsiveness of the brain stem respiratory centers to both increases in carbon dioxide tension and electrical stimulation.

Pentazocine causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origins may produce similar findings). Marked mydriasis rather than miosis may be seen due to hypoxia in overdose situations.

# Effects on the Gastrointestinal Tract and Other Smooth Muscle

Pentazocine causes a reduction in motility associated with an increase in smooth muscle tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone may be increased to the point of spasm, resulting in constipation. Other opioid- induced effects may include a reduction in biliary and ncreatic secretions, spasm of sphincter of Oddi, and transient elevations in serum amylase.

# Effects on the Cardiovascular System

Pentazocine produces peripheral vasodilation which may result in orthostatic hypotension or syncope Manifestations of histamine release and/or peripheral vasodilation may include pruritus, flushing, red eyes sweating, and/or orthostatic hypotension

# Effects on the Endocrine System

Opioids inhibit the secretion of adrenocorticotropic hormone (ACTH), cortisol, and luteinizing hormone (LH) in humans [see **Adverse Reactions**]. They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon.

Chronic use of opioids may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The causal role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date [see **Adverse Reactions**].

### Effects on the Immune System

Opioids have been shown to have a variety of effects on components of the immune system. The clinical significance of these findings is unknown. Overall, the effects of opioids appear to be modestly

### Concentration-Efficacy Relationships

The minimum effective analgesic concentration will vary widely among patients, especially among patients who have been previously treated with potent agonist opioids. The minimum effective analgesic concentratior of pentazocine for any individual patient may increase over time due to an increase in pain, the developmen of a new pain syndrome, and/or the development of analgesic tolerance [see Dosage and Administration]

### Concentration-Adverse Reaction Relationships

There is a relationship between increasing pentazocine plasma concentration and increasing frequency of dose-related opioid adverse reactions such as nausea, vomiting, CNS effects, and respiratory depression In opioid-tolerant patients, the situation may be altered by the development of tolerance to opioid-related adverse reactions [see Dosage and Administration

### Opioid Antagonist Effects

Pentazocine weakly antagonizes the analgesic effects of morphine, meperidine, and phenazocine; in addition it produces incomplete reversal of cardiovascular, respiratory, and behavioral depression induced by phine and meperidine. Pentazocine has about 1/50 the antagonistic activity of nalorphine. It also has

Naloxone when administered orally at 0.5 mg has no pharmacologic activity. Naloxone hydrochloride administered parenterally at the same dose is an antagonist to pentazocine and a pure antagonist to narcotic analgesics.

Pentazocine and Naloxone Tablets are a potent analgesic when administered orally. However, the presence of naloxone in Pentazocine and Naloxone Tablets is intended to prevent the effect of pentazocine if the product is misused by injection.

Studies in animals indicate that the presence of naloxone does not affect pentazocine analgesia when the combination is given orally. If the combination is given by injection the action of pentazocine is neutralized

## Pharmacokinetics

Onset of significant analgesia usually occurs between 15 and 30 minutes after oral administration, and duration of action is usually three hours or longer.

Pentazocine is well absorbed from the gastrointestinal tract. Concentrations in plasma coincide closely with the onset, duration, and intensity of analgesia. The time to mean peak concentration in 24 normal volunteers was 1.7 hours (range 0.5 to 4 hours) after oral administration and the mean plasma elimination half-life was 3.6 hours (range 1.5 to 10 hours).

Pentazocine is metabolized in the liver and excreted primarily in the urine. The products of the oxidation of the terminal methyl groups and glucuronide conjugates are excreted by the kidney. Eliminatic approximately 60% of the total dose occurs within 24 hours. Pentazocine passes into the fetal circulal

### INDICATIONS AND USAGE entazocine and Naloxone Tablets are indicated for the management of pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate

Because of the risks of addiction, abuse, and misuse, with opioids, even at recommended doses, reserve Pentazocine and Naloxone Tablets for use in patients for whom alternative treatment options [e.g., non-opioid analgesics]

Have not been tolerated, or are not expected to be tolerated Have not provided adequate analgesia, or are not expected to provide adequate analgesia

## CONTRAINDICATIONS

Pentazocine and Naloxone Tablets are contraindicated in patients with:

- Significant respiratory depression [see WARNINGS]
- Significant respiratory depression [See Wannings]
  Acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment [see WARNINGS]. Patients with known or suspected gastrointestinal obstruction, including paralytic ileus [see WARNINGS] Patients with hypersensitivity to either pentazocine, naloxone, or any of the formulation excipients
- (e.g., anaphylaxis) [see WARNINGS].

## WARNINGS

WANNINGS Addiction, Abuse, and Misuse Pentazocine and Naloxone Tablets contain pentazocine, a Schedule IV controlled substance. As an or azocine and Naloxone Tablets expose users to the risks of addiction, abuse, and misuse [see **DRUG ABUSE** AND DEPENDENCE].

Although the risk of addiction in any individual is unknown, it can occur in patients appropriately prescribed Pentazocine and Naloxone Tablets. Addiction can occur at recommended dosages and if the drug is misused or abused.

Assess each patient's risk for opioid addiction, abuse, or misuse prior to prescribing Pentazocine and Naloxone Tablets, and monitor all patients receiving Pentazocine and Naloxone Tablets for the development of these behaviors and conditions. Risks are increased in patients with a personal or family history of substance abuse (including drug or alcohol abuse or addiction) or mental illness (e.g., major depression). The potential for these risks should not, however, prevent the proper management of pain in any given patient. Patients at increased risk may be prescribed opioids such as Pentazocine and Naloxone Tab use in such patients necessitates intensive counseling about the risks and proper use of Pentazocine and Naloxone Tablets along with intensive monitoring for signs of addiction, abuse, and misuse

Opioids are sought by drug abusers and people with addiction disorders and are subject to criminal diversion Consider these risks when prescribing or dispensing Pentazocine and Naloxone Tablets. Strategies to reduce these risks include prescribing the drug in the smallest appropriate quantity and advising the patient on the proper disposal of unused drug [see **PRECAUTIONS**; **Information for Patients**]. Contact local state ofessional licensing board or state controlled substances authority for information on how to prevent and tect abuse or diversion of this product.

# Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a Risk Evaluation and Mitigation Strategy (REMS) for these products. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare providers are strongly encouraged to do all of the following:

• Complete a REMS-compliant education program offered by an accredited provider of continuing education

(CE) or another education program that includes all the elements of the FDA Education Blueprint for Health Care Providers Involved in the Management or Support of Patients with Pain.

Discuss the safe use, serious risks, and proper storage and disposal of opioid analgesics with patients and/or their caregivers every time these medicines are prescribed. The Patient Counseling Guide (PCG) can be obtained at this link: http://www.fda.gov/OpioidAnalgesicREMSPCG.
 Emphasize to patients and their caregivers the importance of reading the Medication Guide that they will

receive from their pharmacist every time an opioid analgesic is dispensed to them Consider using other tools to improve patient, household, and community safety, such as patient-prescriber agreements that reinforce patient-prescriber responsibilities.

To obtain further information on the opioid analgesic REMS and for a list of accredited REMS CME/CE, call 800-503-0784, or log on to www.opioidanalgesicrems.com. The FDA Blueprint can be found at www.fda.gov/OpioidAnalgesicREMSBlueprint.

# Life-Threatening Respiratory Depression

Serious, life-threatening nor fatal respiratory depression has been reported with the use of opioids, even when used as recommended. Respiratory depression, if not immediately recognized and treated, may lead to respiratory arrest and death. Management of respiratory depression may include close observation, asures, and use of opioid antagonists, depending on the patient's clinical status [see OVERDOSAGE]. Carbon dioxide (CO<sub>2</sub>) retention from opioid-induced respiratory depression can exacerba he sedating effects of opioids.

While serious, life-threatening, or fatal respiratory depression can occur at any time during the use of Pentazocine and Naloxone Tablets, the risk is greatest during the initiation of therapy or following a dosage increase. Monitor patients closely for respiratory depression, especially within the first 24 to 72 hours of nitiating therapy with and following dosage increases of Pentazocine and Naloxone Tablets

To reduce the risk of respiratory depression, proper dosing and titration of Pentazocine and Naloxone Tablets are essential [see **DOSAGE AND ADMINISTRATION**]. Overestimating the Pentazocine and Naloxone Tablets dosage when converting patients from another opioid product can result in a fatal overdose with the first dose Accidental ingestion of even one dose of Pentazocine and Naloxone Tablets, especially by children, can result

nestinating state of the control of Administration].

Neonatal Opioid Withdrawal Syndrome
Prolonged use of Pentazocine and Naloxone Tablets during pregnancy can result in withdrawal in the neonate.
Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life-threatening in not recognized and treated, and requires management according to protocols developed neonatology experts. Observe newborns for signs of neonatal opioid withdrawal syndrome and manage accordingly. Advise pregnant women using opioids for a prolonged period of the risk of neonatal opioid withdrawa syndrome and ensure that appropriate treatment will be available [see PRECAUTIONS; Information for

## Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants

Profound sedation, respiratory depression, coma, and death may result from the concomitant use of Pentazorine and Naloxone Tablets with benzodiazepines or other CNS depressants (e.g., non-benzodiazepine sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, other opioids, alcohol). Because of these risks, reserve concomitant prescribing of these drugs for use in patients or whom alternative treatment options are inadequate.

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazen observational studies have definished with a concentration use of opinion analysis and behaviorable to creases the risk of drug-related mortality compared to use of opinion analysis alone. Because of similar harmacological properties, it is reasonable to expect similar risk with the concomitant use of other CNS epressant drugs with opinion analysis (see PRECAUTIONS, Drug Interactions).

depressant drugs with opinion angestes (see "herbor orbot, bright metactions).

If the decision is made to prescribe a benzodiazepine or other CNS depressant concomitantly with an opioid analgesic, prescribe the lowest effective dosages and minimum durations of concomitant use. In patients already receiving an opioid analgesic, prescribe a lower initial dose of the benzodiazepine or other CNS depressant than indicated in the absence of an opioid, and titrate based on clinical response. If an opioid analgesic is initiated in a patient already taking a benzodiazepine or other CNS depressant, prescribe a lower initial dose of the opioid analgesic, and titrate based on clinical response. Follow patients closely for signs and symptoms of respiratory depression and sedation

Advise both patients and caregivers about the risks of respiratory depression and sedation when Pentazocine Advise both patients and caregivers about the risks of respiratory depression and sedation when Pentazotine and Naloxone Tablets is used with benzodiazepines or other CNS depressants (including alcohal dillicit drugs). Advise patients not to drive or operate heavy machinery until the effects of concomitant use of the benzodiazepine or other CNS depressant have been determined. Screen patients for risk of substance use disorders, including opioid abuse and misuse, and warn them of the risk for overdose and death associated vith the use of additional CNS depressants including alcohol and illicit drugs [see PRECAUTIONS: Information for Patients, Drug Interactions]

## Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly, Cachectic, or Debilitated Patients The use of Pentazocine and Naloxone Tablets in patients with acute or severe bronchial asthma in an

inmonitored setting or in the absence of resuscitative equipment is contraindicated.

Patients with Chronic Pulmonary Disease: Pentazocine and naloxone-treated patients with significant chronic radients with Chronic Pullinoiry Disease, reliazochie and haloxone-treated patients with significant obstructive pulmonary disease or cor pulmonale, and those with a substantially decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory depression are at increased risk of decreased espiratory drive including apnea, even at recommended dosages of Pentazocine and Naloxone Tablets [see

Elderly, Cachectic, or Debilitated Patients: Life-threatening respiratory depression is more likely to occur in elderly, cachectic, or debilitated patients because they may have altered pharmacokinetics or altered clearance elderly, cachectic, or debilitated patients because they may have compared to younger, healthier patients [see WARNINGS].

Monitor such patients closely, particularly when initiating and titrating Pentazocine and Naloxone Tablets and when Pentazocine and Naloxone Tablets are given concomitantly with other drugs that depress respiration [see WARNINGS]. Alternatively, consider the use of non-opioid analgesics in these patients.

Cases of adrenal insufficiency have been reported with opioid use, more often following greater than 1 month of use. Presentation of adrenal insufficiency may include non-specific symptoms and signs including nausea comitting, anorexia, fatigue, weakness, dizziness, and low blood pressure. If adrenal insufficiency is suspected, confirm the diagnosis with diagnostic testing as soon as possible. If adrenal insufficiency is diagnosed, treat with physiologic replacement doses of corticosteroids. Wean the patient off of the opioid to allow adrenal function to recover and continue corticosteroid treatment until adrenal function recovers. Other opioids may be tried as some cases reported use of a different opioid without recurrence of adrena nsufficiency. The information available does not identify any particular opioids as being more likely to be ssociated with adrenal insufficiency

### Severe Hypotension

Pentazocine and Naloxone Tablets may cause severe hypotension including orthostatic hypotension and syncope in ambulatory patients. There is increased risk in patients whose ability to maintain blood pressure syncope in almountly patients. There is included that in patients winds admin to find pressure has already been compromised by a reduced blood volume or concurrent administration of certain CNS depressant drugs (e.g., phenothiazines or general anesthetics) [see PRECAUTIONS; Information for Patients]. Monitor these patients for signs of hypotension after initiating or titrating the dosage of Pentazocine and Naloxone Tablets. In patients with circulatory shock, Pentazocine and Naloxone Tablets may cause vasodilation that can further reduce cardiac output and blood pressure. Avoid the use of Pentazocine and Naloxone Tablets in patients with circulatory shock.

# Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired

n patients who may be susceptible to the intracranial effects of CO<sub>2</sub> retention (e.g., those with evidence of ncreased intracranial pressure or brain tumors), Pentazocine and Naloxone Tablets may reduce respiratory drive, and the resultant CO<sub>2</sub> retention can further increase intracranial pressure. Monitor such patients for signs of sedation and respiratory depression, particularly when initiating therapy with Pentar

Opioids may also obscure the clinical course in a patient with a head injury. Avoid the use of Pentazocine and Naloxone Tablets in patients with impaired consciousness or coma

# Risks of Use in Patients with Gastrointestinal Conditions

Pentazocine and Naloxone Tablets are contraindicated in patients with known or suspected gastrointestinal obstruction, including paralytic ileus.

The administration of Pentazocine and Naloxone Tablets or other opioids may obscure the diagnosis or clinical course in patients with acute abdominal conditions. Pentazocine and Naloxone Tablets may cause spasm of the sphincter of Oddi. Opioids may cause increases in serum amylase. Monitor patients with biliary tract disease, including acute pancreatitis, for worsening

# symptoms.

Withdrawal Do not abruptly discontinue Pentazocine and Naloxone Tablets in a patient physically dependent on opioids. When discontinuing Pentazocine and Naloxone Tablets in a physically dependent patient, gradually taper the dosage. Rapid tapering of Pentazocine and Naloxone Tablets in a physically dependent patient, gradually taper the dosage. Rapid tapering of Pentazocine and Naloxone Tablets in a patient physically dependent on opioids may lead to a withdrawal syndrome and return of pain [see **Dosage and Administration, Drug Abuse and** 

Additionally, the use of Pentazocine and Naloxone Tablets, a mixed agonist/antagonist opioid analgesic patients who are receiving a full opioid agonist analgesic may reduce the analgesic effect and/or precipitate withdrawal symptoms. Avoid concomitant use of Pentazocine and Naloxone Tablets with a full opioid agonist analgesic.

# **Acute CNS Manifestations**

Patients receiving therapeutic doses of Pentazocine and Naloxone Tablets have experienced hallucinations (usually visual), disorientation, and confusion which have cleared spontaneously within a period of hours. The mechanism of this reaction is not known. Such patients should be very closely observed and vital signs checked. If the drug is reinstituted, it should be done with caution since these acute CNS manifestations

taken orally and will not interfere with the pharmacologic action of pentazocine. However, this amount of naloxone given by injection has profound antagonistic action to narcotic analgesics. Severe, even lethal, consequences may result from misuse of tablets by injection either alone or in

The amount of naloxone present in Pentazocine and Naloxone Tablets (0.5 mg per tablet) has no action when

combination with other substances, such as pulmonary emboli, vascular occlusion, ulceration and abscesses and withdrawal symptoms in narcotic dependent individuals.

### Increased Risk of Seizures in Patients with Seizure Disorders The pentazocine in Pentazocine and Naloxone Tablets may increase the frequency of seizures in patients

with seizure disorders, and may increase the risk of seizures occurring in other clinical settings associated with seizures. Monitor patients with a history of seizure disorders for worsened seizure control during Pentazocine and Naloxone Tablets therapy

Risks of Driving and Operating Machinery
Pentazocine and Naloxone Tablets may impair the mental or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery. Warn patients not to drive or operate dangerous machinery unless they are tolerant to the effects of Pentazocine and Naloxone Tablets and know how they will react to the medication [see Patient Counseling Information].

Chronic use of opioids may cause reduced fertility in females and males of reproductive potential. It is not known whether these effects on fertility are reversible [see ADVERSE REACTIONS]. PRECAUTIONS

## Porphyria

Particular caution should be exercised in administering pentazocine to patients with porphyria since it may provoke an acute attack in susceptible individuals.

## Cardiovascular Disease

have nausea or vomiting.

Pentazocine can elevate blood pressure, possibly through the release of endogenous catecholamines. Particular caution should be exercised in conditions where alterations in vascular resistance and blood pressure might be particularly undesirable, such as in the acute phase of myocardial infarction. Pentazocine and Naloxone Tablets should be used with caution in patients with myocardial infarction who

### Impaired Renal or Hepatic Function

Decreased metabolism of pentazocine by the liver in extensive liver disease may predispose to accentuation of side effects. Although laboratory tests have not indicated that pentazocine causes or increases renal or hepatic impairment, the drug should be administered with caution to patients with such impairment.

Narcotic drug products are generally considered to elevate biliary tract pressure for varying periods following dministration. Some evidence suggests that pentazocine may differ from other marketed narcotics in spect (i.e., it causes little or no elevation in biliary tract pressures). The clinical significance of these

### Information for Patients

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Storage and Disposal
Because of the risks associated with accidental ingestion, misuse, and abuse, advise patients to store
Pentazocine and Naloxone Tablets securely, out of sight and reach of children, and in a location not accessible
by others, including visitors to the home [see WARNINGS, DRUG ABUSE AND DEPENDENCE]. Inform
patients that leaving Pentazocine and Naloxone Tablets unsecured can pose a deadly risk to others in the

Advise patients and caregivers that when medicines are no longer needed, they should be disposed of promptly. Expired, unwanted, or unused Pentazocine and Naloxone Tablets should be disposed of by flushing the unused medication down the toilet if a drug take-back option is not readily available. Inform patients that they can visit www.fda.gov/drugdisposal for a complete list of medicines recommended for disposal by flushing, as well as additional information on disposal of unused medicines

Addiction, Abuse, and Misuse Inform patients that the use of Pentazocine and Naloxone Tablets, even when taken as recommended, can result in addiction, abuse, and misuse, which can lead to overdose and death [see WARNINGS]. Instruct patients not to share Pentazocine and Naloxone Tablets with others and to take steps to protect Pentazocine and Naloxone Tablets from theft or misuse. Life-Threatening Respiratory Depression

# Inform patients of the risk of life-threatening respiratory depression, including information that the risk is greatest when starting Pentazocine and Naloxone Tablets or when the dosage is increased, and that it can occur even at recommended dosages [see WARNINGS]. Advise patients how to recognize respiratory depression and to seek medical attention if breathing difficulties develop

Accidental Ingestion
Inform patients that accidental ingestion, especially by children, may result in respiratory depression or

# death [see WARNINGS].

Interactions with Benzodiazepines and Other CNS Depressants
Inform patients and caregivers that potentially fatal additive effects may occur if Pentazocine and Naloxone
Tablets are used with benzodiazepines or other CNS depressants, including alcohol, and not to use these
drugs concomitantly unless supervised by a healthcare provider [see WARNINGS, PRECAUTIONS; Drug Interactions].

# Serotonin Syndrom

Inform patients that opioids could cause a rare but potentially life-threatening condition resulting from concomitant administration of serotonergic drugs. Warn patients of the symptoms of serotonin syndrome and to seek medical attention right away if symptoms develop. Instruct patients to inform the relatthcare provider if they are taking, or plan to take serotonergic medications [see PRECAUTIONS; Drug Interactions]. Adrenal Insufficiency ients that opioids could cause adrenal insufficiency, a potentially life threatening condition. Adrenal

# insufficiency may present with non-specific symptoms and signs such as nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. Advise patients to seek medical attention if they experience a constellation of these symptoms [see **WARNINGS**]. Important Administration Instructions

Advise patients now to properly take Pentazocine and Naloxone Tablets.

Advise patients not to adjust the dose of Pentazocine and Naloxone Tablets without consulting with a physician or other healthcare professional.

If patients have been receiving treatment with Pentazocine and Naloxone Tablets for more than a few weeks and cessation of therapy is indicated, counsel them on the importance of safely tapering the dose as abruptly discontinuation of the medication could precipitate withdrawal symptoms. Provide a dose schedule to accomplish a gradual discontinuation of the medication. [see DOSAGE AND

# ADMINISTRATION] Important Discontinuation Instructions In order to avoid developing withdrawal symptoms, instruct patients not to discontinue Pentazocine and

Naloxone Tablets without first discussing a tapering plan with the prescriber [see DOSAGE AND ADMINISTRATION nform patients that Pentazocine and Naloxone Tablets may cause orthostatic hypotension and syncope.

Instruct patients how to recognize symptoms of low blood pressure and how to reduce the risk of serious

### ences should hypotension occur (e.g., sit or lie down, carefully rise from a sitting or lying position) see WARNINGS

**Anaphylaxis** rm patients that anaphylaxis have been reported with ingredients contained in Pentazocine and Naloxone Tablets. Advise patients how to recognize such a reaction and when to seek medical attention *[see* Contraindications, Adverse Reactions]

# Inform female patients of reproductive potential that prolonged use of Pentazocine and Naloxone Tablets

during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not ognized and treated [see WARNINGS, PRECAUTIONS; Pregnancy] Embryo-Fetal Toxicity Inform female patients of reproductive potential that Pentazocine and Naloxone Tablets can cause fetal

harm and to inform the healthcare provider of a known or suspected pregnancy [see PRECAUTIONS;

Lactation
Advise nursing mothers to monitor infants for increased sleepiness (more than usual), breathing difficulties, or limpness. Instruct nursing mothers to seek immediate medical care if they notice these signs [see


<u>Driving or Operating Heavy Machinery</u> Inform patients that Pentazocine and Naloxone Tablets may impair the ability to perform potentially hazardous activities such as driving a car or operating heavy machinery. Advise patients not to perform such tasks until they know how they will react to the medication [see **PRECAUTIONS**].

# Advise patients of the potential for severe constipation, including management instructions and when to seek medical attention (see **ADVERSE REACTIONS, CLINICAL PHARMACOLOGY**). **Drug Interactions**

Benzodiazenines and Other Central Nervous System (CNS) Depressants

Due to additive pharmacologic effect, the concomitant use of benzodiazepines or other CNS depressants including alcohol, benzodiazepines and other sedative hypnotics, anxiolytics, and tranquilizers, muscle relaxants, general anesthetics, antipsychotics, and other opioids, can increase the risk of hypotension, respiratory depression, profound sedation, coma, and death

CONTROL Proof Date: 08/22/2019 Proof Time: 02:11 PM Prepared by: suzanne NP Item#: NOVE-NP\_41212 Size: 12.25 x 16.25 (folded: 1.25 x 1.25) Type size: 6 pt. Item Iss./Rev. Date: Revised: 08/2019 Cust. Part No.: PI6800000211 PO No.: Description: Pentazocine Hcl & Naloxone Hcl Tabs, Pl6800000211 (Lup Label: Lupin Customer: Novel Bar code details: Type: UPC-A Code: 43386-680-01 Notes: ■ Approved Resubmit Signature: Date:



**Naloxone Tablets** 

USP

08/22/2019 02:11 PM / NP Item# NOVE-NP\_41212 / page 1 of 2



Reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate. Limit dosages and durations to the minimum required. Follow patients closely for signs of respiratory depression and sedation [see **WARNINGS**].

### Serotonergic Drugs

The concomitant use of opioids with other drugs that affect the serotonergic neurotransmitter system, such The concomitant use of options with other drugs that affect the serotonergic neurotransmitter system, such as selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), triptans, 5-HT3 receptor antagonists, drugs that affect the serotonin neurotransmitter system (e.g., mirtazapine, trazodone, tramadol), certain muscle relaxants (i.e., cyclobenzaprine, metaxalone), monoamine oxidase (MAO) inhibitors (those intended to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue), has resulted in serotonin syndrome. [see PRECAUTIONS; Information for Patients]

If concomitant use is warranted, carefully observe the patient, particularly during treatment initiation and dose adjustment. Discontinue Pentazocine and Naloxone Tablets if serotonin syndrome is suspected.

Monoamine Oxidase Inhibitors (MAOIs)
Concomitant use of monoamine oxidase inhibitors (MAOIs) with Pentazocine and Naloxone Tablets may cause CNS excitation and hypertension through their respective effects on catecholamines. Caution should therefore be observed in administering Pentazocine and Naloxone Tablets to patients who are currently receiving MAOIs or who have received them within the preceding 14 days.

Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics
Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics such as butorphanol, nalbuphine, pentazocine, buprenorphine, may reduce the analgesic effect of Pentazocine and Naloxone Tablets and/or precipitate withdrawal symptoms.

Avoid concomitant use of these drugs.

### Muscle Relaxants

The Concomitant use of opioids and muscle relaxants may enhance the neuromuscular blocking action of skeletal muscle relaxants and produce an increased degree of respiratory depression.

Monitor patients for signs of respiratory depression that may be greater than otherwise expected and decrease the dosage of Pentazocine and Naloxone Tablets and/or the muscle relaxant as necessary

### Diuretics

Opioids can reduce the efficacy of diuretics by inducing the release of antidiuretic hormone

# Monitor patients for signs of diminished diuresis and/or effects on blood pressure and increase the dosage

of the digretic as needed

Anticholinergic Drugs
The concomitant use of anticholinergic drugs may increase risk of urinary retention and/or severe stipation, which may lead to paralytic ileus.

Monitor patients for signs of urinary retention or reduced gastric motility when Pentazocine and Naloxone Tablets is used concomitantly with anticholinergic drugs.

Todacco Smoking tobacco could enhance the metabolic clearance rate of pentazocine reducing the clinical effectiveness of a standard dose of pentazocine.

## Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term animal studies have not been completed to evaluate the carcinogenic potential of the combination or individual components of Pentazocine and Naloxone Tablets.

Studies to evaluate the mutagenic potential of the components of Pentazocine and Naloxone Tablets have not been conducted.

### Impairment of Fertility

Studies in animals to evaluate the impact of Pentazocine and Naloxone Tablets on fertility have not been

The daily administration of 4 mg/kg to 20 mg/kg pentazocine subcutaneously to female rats during a 14 day pre-mating period and until the 13th day of pregnancy did not have any adverse effects on the fertility rate.

## Prennancy

## Risk Summary

Prolonged use of opioid analgesics during pregnancy can cause neonatal opioid withdrawal syndrome [see Warnings and Precautions (5.3)]. There are no available data with Pentazocine and Naloxone Tablets in pregnant women to inform a drug-associated risk for major birth defects and miscarriage. In animal reproduction studies, pentazocine administered subcutaneously to pregnant hamsters during the early gestational period produced neural tube defects (i.e., exencephaly and cranioschisis) at 2.6 times the maximum daily dose (MDD). In pregnant rats administered pentazocine:naloxone during organogenesis, there were increased incidences of resorptions and extra ribs at 0.2 times the MDD. There was no evidence of malformations in rats or rabbits [see Data]. Based on animal data, advise pregnant women of the potential risk to a fetus. The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Prolonged use of opioid analgesics during pregnancy for medical or nonmedical purposes can result in physical dependence in the neonate and neonatal opioid withdrawal syndrome shortly after birth.

Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high pitched cry, tremor, vomiting, diarrhea and failure to gain weight. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by the newborn. Observe newborns for symptoms of neonatal opioid withdrawal syndrome and manage accordingly [see WARNINGS].

## Labor or Delivery

Opioids cross the placenta and may produce respiratory depression and psycho-physiologic effects in neonates. An opioid antagonist, such as naloxone, must be available for reversal of opioid-induced respiratory depression in the neonate. Pentazocine and Naloxone Tablets are not recommended for use in pregnant women during or immediately prior to labor, when other analgesic techniques are more appropriate. Opioid analgesics, including Pentazocine and Naloxone Tablets, can prolong labor through actions which temporarily reduce the strength, duration, and frequency of uterine contractions. However, this effect is not consistent and may be offset by an increased rate of cervical dilation, which tends to shorten labor. Monitor neonates exposed to opioid analgesics during labor for signs of excess sedation and respiratory depression

Affilm Data I or a published report, a single dose of pentazocine administered to pregnant hamsters on Gestation Day 8 increased the incidence of neural tube defects (exencephaly and cranioschisis) at a dose of 196 mg/kg, SC (2.6-times the maximum daily human dose (MDD) of 600 mg/day pentazocine (12 tablets) on a mg/m² basis). No evidence of neural tube defects were reported following a dose of 98 mg/kg (1.3 times the MDD).

Animal reproduction studies testing the combination of pentazocine and naloxone during organogenesis have been completed in rats and rabbits. In rats, a pentazocine:naloxone dose of 64 mg/kg:0.64 mg/kg via oral gavage from Gestation Day 6 to 15 increased the incidences of resorptions and extra ribs (0.2 times the maximum daily human dose of pentazocine via 12 tablets on a mg/m² basis). There were no clear treatment related effects in rabbits treated from Gestation Day 6 to 18 with a pentazocine:naloxone dose of up to 64 mg/kg:0.64 mg/kg via oral gavage (0.3-times the maximum daily human dose of pentazocine via 12 tablets on a mg/m² basis). on a mg/m2 basis

## Lactation

rtazocine is excreted in human milk. Caution should be exercised when Pentazocine and Naloxone Tablets are administered to a nursing woman.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Pentazocine and Naloxone Tablets and any potential adverse effects on the breastfed infant from Pentazocine and Naloxone Tablets or from the underlying maternal condition.

## Clinical Considerations

nts exposed to pentazocine and naloxone through breast milk should be monitored for excess sedation and respiratory depression. Withdrawal symptoms can occur in breastfed infants when maternal administration of an opioid analogoic is stopped, or when breast-feeding is stopped.

## Pediatric Use

Safety and effectiveness in pediatric patients below the age of 12 years have not been established.

## Geriatric Use

Geriamic use Elderly patients (aged 65 years or older) may have increased sensitivity to Pentazocine and Naloxone Tablets. In general, use caution when selecting a dosage for an elderly patient, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

Respiratory depression is the chief risk for elderly patients treated with opioids, and has occurred after large initial doses were administered to patients who were not opioid-tolerant or when opioids were co-administered with other agents that depress respiration. Titrate the dosage of Pentazocine and Naloxone Tablets slowly in geriatric patients [see WARNINGS].

Pentazocine and Naloxone are known to be substantially excreted by the kidney, and the risk of adverse eactions to this drug may be greater in patients with impaired renal function. Because elderly patients are nore likely to have decreased renal function, care should be taken in dose selection, and it may be useful to

### ADVERSE REACTIONS

The following adverse reactions associated with the use of Pentazocine and Naloxone were identified in clinical studies or postmarketing reports. Because some of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Cardiovascular - Hypertension, hypotension, circulatory depression, tachycardia, syncope

Respiratory - Rarely, respiratory depression.

Acute CNS Manifestations - Hallucinations (usually visual), disorientation, and confusion.

Other CNS Effects - Grand mal convulsions, increase in intracranial pressure, dizziness, lightheadedness ons, sedation, euphoria, headache, confusion, disprientation; infrequently weakness, disturbed dreams, insomnia, syncope, and depression; and rarely tremor, irritability, excitement, tinnitus

Autonomic - Sweating; infrequently flushing; and rarely chills.

Gastrointestinal - Nausea, vomiting, constipation, diarrhea, anorexia, dry mouth, biliary tract spasm, and

**Allergic** - Edema of the face; anaphylactic shock; dermatitis, including pruritus; flushed skin, including plethora; infrequently rash, and rarely urticaria.

Ophthalmic - Visual blurring and focusing difficulty, miosis.

Hematologic - Depression of white blood cells (especially granulocytes), with rare cases of agranulocytosis, which is usually reversible, moderate transient eosinophilia

# Dependence and Withdrawal Symptoms - (See WARNINGS, PRECAUTIONS, and DRUG ABUSE AND

**Other -** Urinary retention, paresthesia, serious skin reactions, including erythema multiforme, Stevens-Johnson syndrome toxic epidermal necrolysis, and alterations in rate or strength of uterine contractions during labor.

- <u>Serotonin syndrome</u>; Cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of opioids with serotonergic drugs.
- Adrenal insufficiency: Cases of adrenal insufficiency have been reported with opioid use, more often
- <u>Anaphylaxis:</u> Anaphylaxis has been reported with ingredients contained in Pentazocine and Naloxone
  Tablete
  Table
- Androgen deficiency: Cases of androgen deficiency have occurred with chronic use of opioids [see Clinical Pharmacoloov].

## DRUG ABUSE AND DEPENDENCE

Pentazocine and Naloxone Tablets contain pentazocine, a Schedule IV controlled substance

Pentazocine and Naloxone Tablets contain pentazocine, a substance with a high potential for abuse similar to other opioids including tramadol. Pentazocine and Naloxone Tablets can be abused and is subject to misuse, addiction, and criminal diversion [see **WARNINGS**].

All patients treated with opioids require careful monitoring for signs of abuse and addiction, because use of opioid analgesic products carries the risk of addiction even under appropriate medical use.

Prescription drug abuse is the intentional non-therapeutic use of a prescription drug, even once, for its

Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that develop after repeated substance use and includes: a strong desire to take the drug, difficulties in controlling its use, persisting in its use despite harmful consequences, a higher priority given to drug use than to other activities and obligations, increased tolerance, and sometimes a physical withdrawal.

"Drug-seeking" behavior is very common in persons with substance use disorders. Drug-seeking factics "Drug-seeking" behavior is very common in persons with substance use disorders. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing, or referral, repeated "loss" of prescriptions, tampering with prescriptions, and reluctance to provide prior medical records or contact information for other treating health care provider(s). "Doctor shopping" (visiting multiple prescribers to obtain additional prescriptions) is common among drug abusers and people suffering from untreated addiction. Preoccupation with achieving adequate pain relief can be appropriate behavior in a patient with poor pain control.

Abuse and addiction are separate and distinct from physical dependence and tolerance. Healthcare providers should be aware that addiction may not be accompanied by concurrent tolerance and symptoms of physical dependence in all addicts. In addition, abuse of opioids can occur in the absence of true addiction.

Pentazocine and Naloxone Tablets, like other opioids, can be diverted for non-medical use into illicit channels of distribution. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests, as required by state and federal law, is strongly advised.

Proper assessment of the patient, proper prescribing practices, periodic re-evaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs

## Risks Specific to Abuse of Pentazocine and Naloxone Tablets

Pentazocine and Naloxone Tablets is for oral use only. Abuse of Pentazocine and Naloxone Tablets poses a reinazoonie and valoxone rabiets is for oral use only. Aduse of Penazoonie and valoxone rabiets poses in risk of overdose and death. The risk is increased with concurrent use of Pentazocine and Naloxone Tablets with alcohol and other central nervous system depressants. Parenteral drug abuse is commonly associated with transmission of infectious diseases such as hepatitis and HIV.

Both tolerance and physical dependence can develop during chronic opioid therapy. Tolerance is the need for increasing doses of opioids to maintain a defined effect such as analgesia (in the absence of disease progression or other external factors). Tolerance may occur to both the desired and undesired effects of drugs, and may develop at different rates for different effects.

hysical dependence is a physiological state in which the body adapts to the drug after a period of regular exposure, resulting in withdrawal symptoms after abrupt discontinuation or a significant dosage reduction of a drug. Withdrawal also may be precipitated through the administration of drugs with opioid antagonist activity (e.g., naloxone, nalmefene), mixed agonist/antagonist analgesics (e.g., butorphanol, nalbuphine), or partial agonists (e.g., buprenorphine). Physical dependence may not occur to a clinically significant degree until after several days to weeks of continued opioid usage.

Do not abruptly discontinue Pentazocine and Naloxone Tablets in a patient physically dependent on opioids. Rapid tappring of Pentazocine and Naloxone Tablets in a patient physically dependent on opioids may lead to serious withdrawal symptoms, uncontrolled pain, and suicide. Rapid discontinuation has also been associated with attempts to find other sources of opioid analgesics, which may be confused with

When discontinuing Pentazocine and Naloxone Tablets, gradually taper the dosage using a natientspecific When discontinuing Pentazocine and Naloxone lablets, gradually taper the dosage using a patientspecii plan that considers the following: the dose of Pentazocine and Naloxone Tablets the patient be been takin the duration of treatment, and the physical and psychological attributes of the patient. To improve the likelihood of a successful taper and minimize withdrawal symptoms, it is important that the opioid taper is schedule is agreed upon by the patient. In patients taking opioids for a long duration at high doses, ensu that a multimodal approach to pain management, including mental health support (if needed), is in place pri to initiating an opioid analgesic taper [see DOSAGE AND ADMINISTRATION, WARNINGS].

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal signs [see PRECAUTIONS; Pregnancy]

<u>Clinical Presentation</u> Acute overdose with Pentazocine and Naloxone Tablets can be manifested by respiratory depression, somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and, in some cases, pulmonary edema, bradycardia, hypotension, partial or complete airway obstruction, atypical snoring, and death. Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations.

For pentazocine alone in single doses above 60 mg there have been reports of the occurrence of rphine-like psychotomimetic effects such as anxiety, nightmares, strange thoughts, and hallucinations. nnolence, marked respiratory depression associated with hypertension and tachycardia have also resulted as have seizures, hypotension, dizziness, nausea, vomiting, lethargy, and paresthesias. The respiratory depression is antagonized by naloxone (see Treatment). Circulatory failure and deepening coma may occur in more severe cases, particularly in patients who have also ingested other CNS depressants such as alcohol, edative/hypnotics, or antihistan

## Treatment of Overdose

case of overdose, priorities are the reestablishment of a patent and protected airway and institution of sisted or controlled ventilation, if needed. Employ other supportive measures (including oxygen and soppressors) in the management of circulatory shock and pulmonary edema as indicated. Cardiac arrest arrhythmias will require advanced life-support techniques.

The opioid antagonist, naloxone or nalmefene, are specific antidotes to respiratory depression resulting from opioid overdose. For clinically significant respiratory or circulatory depression secondary to pentazocine overdose, administer an opioid antagonist. As pentazocine is a mixed opioid agonist/antagonist, larger doses of naloxone or nalmefene may be needed to reverse the effects of an overdose.

In an individual physically dependent on opioids, administration of the recommended usual dosage of the intagonist will precipitate an acute withdrawal syndrome. The severity of the withdrawal symptoms ienced will depend on the degree of physical dependence and the dose of the antagonist admi f a decision is made to treat serious respiratory depression in the physically dependent patient nistration of the antagonist should be begun with care and by titration with smaller than usual doses of the antagonist.

### DOSAGE AND ADMINISTRATION

mportant Dosage and Administration Instructions

Jse the lowest effective dosage for the shortest duration consistent with individual patient treatment goals [see WARNINGS].

Monitor patients closely for respiratory depression, especially within the first 24 to 72 hours of initiating therapy and following dosage increases with Pentazocine and Naloxone Tablets and adjust the dosage accordingly [see WARNINGS].

### Initial Dosage

Use of Pentazocine and Naloxone Tablets as the First Opioid Analgesic

should not exceed 12 tablets.

Conversion from Other Opioids to Pentazocine and Naloxone Tablets
There is inter-patient variability in the potency of opioid drugs and opioid formulations. Therefore, a conservative approach is advised when determining the total daily dosage of Pentazocine and Naloxone Tablets. It is safer to underestimate a patient's 24-hour Pentazocine and Naloxone Tablets dosage than to overestimate the 24-hour Pentazocine and Naloxone Tablets dosage and manage an adverse reaction due to

Titration and Maintenance of Therapy
Individually titrate Pentazocine and Naloxone Tablets to a dose that provides adequate analgesia and minimizes adverse reactions. Continually reevaluate patients receiving Pentazocine and Naloxone Tablets to assess the maintenance of pain control and the relative incidence of adverse reactions, as well as monitoring for the development of addiction, abuse, or misuse [see WARNINGS]. Frequent communication is important among the prescriber, other members of the healthcare team, the patient, and the caregiver/family during periods of changing analgesic requirements, including initial titration.

increasing the Pentazocine and Naloxone Tablets dosage. If unacceptable opioid- related adverse reactions are observed, consider reducing the dosage. Adjust the dosage to obtain an appropriate balance between management of pain and opioid-related adverse reactions.

### Safe Reduction or Discontinuation of Pentazocine and Naloxone Tablets

When a decision has been made to decrease the dose or discontinue therapy in an opioid-dependent patient taking Pentazocine and Naloxone Tablets, there are a variety of factors that should be considered, including the dose of Pentazocine and Naloxone Tablets the patient has been taking, the duration of treatment, the type of pain being treated, and the physical and psychological attributes of the patient. It is important to ensure ongoing care of the patient and to agree on an appropriate tapering schedule and follow-up plan so that patient and provider goals and expectations are clear and realistic. When opioid analgesics are being discontinued due to a suspected substance use disorder, evaluate and treat the patient, or refer for evaluation and treatment of the substance use disorder. Treatment should include evidence-based approaches, such as medication assisted treatment of opioid use disorder. Complex patients with co-morbid pain and substance use disorders may benefit from referral to a specialist.

of time may tolerate a more rapid taper.

It may be necessary to provide the natient with lower dosage strengths to accomplish a successful taper It may be necessary to provide the patient with lower dosage strengths to accomplish a successful taper. Reassess the patient frequently to manage pain and withdrawal symptoms, should they emerge. Common withdrawal symptoms include restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, myalgia, and mydriasis. Other signs and symptoms also may develop, including irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, or increased blood pressure, respiratory rate, or heart rate. If withdrawal symptoms arise, it may be necessary to pause the taper for a period of time or raise the dose of the opioid analgesic to the previous dose, and then proceed with a slower taper. In addition, monitor patients for any changes in mood, emergence of suicidal thoughts, or use of other substances.

When managing patients taking opioid analgesics, particularly those who have been treated for a long duration and/or with high doses for chronic pain, ensure that a multimodal approach to pain management, including mental health support (if needed), is in place prior to initiating an opioid analgesic taper. A multimodal approach to pain management may optimize the treatment of chronic pain, as well as assist with the successful tapering of the opioid analgesic [see WARNINGS/Withdrawal, DRUG ABUSE AND DEPENDENCE!] DEPENDENCE1.

## HOW SUPPLIED

Pentazocine and Naloxone Tablets USP are light vellow, capsule shaped tablets debossed "NL" on left side and "680" on the right side of the bisect and plain on the other side, supplied in bottles of 100 and 500.

Bottles of 100 (NDC 43386-680-01). Bottles of 500 (NDC 43386-680-05).

Store at 20° to 25°C (68° to 77°F) [See LISP Controlled Boom Temperature]

# **Medication Guide**

# Pentazocine and Naloxone (pen taz' oh seen and nal ox' one) Tablets



### Pentazocine and Naloxone Tablets are:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage moderate to severe pain, when other pain treatments such as non-opioid pain
- medicines do not treat your pain well enough or you cannot tolerate them.

  An opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.

### Important information about Pentazocine and Naloxone Tablets:

- Get emergency help right away if you take too many Pentazocine and Naloxone Tablets (overdose). When you first start taking Pentazocine and Naloxone Tablets, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur. Taking Pentazocine and Naloxone Tablets with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your Pentazocine and Naloxone Tablets. They could die from taking it. Selling or giving away Pentazocine and Naloxone Tablets is against the law.
- Store Pentazocine and Naloxone Tablets securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home.

### Do not take Pentazocine and Naloxone Tablets if you have:

- severe asthma, trouble breathing, or other lung problems. a bowel blockage or have parrowing of the stomach or intestines.
- previously had an allergic reaction to pentazocine or naloxone.

# known or suspected gastrointestinal obstruction, including paralytic ileus. Before taking Pentazocine and Naloxone Tablets, tell your healthcare provider if you have

- a history of:
- head injury seizures
   liver kidney thyroid problems problems urinating
   pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.

## Tell your healthcare provider if you are:

- pregnant or planning to become pregnant. Prolonged use of Pentazocine and Naloxone Tablets during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- breastfeeding. Pentazocine and naloxone passes into breast milk and may harm your
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking Pentazocine and Naloxone Tablets with certain other medicines can cause serious side effects that could lead to death.

# When taking Pentazocine and Naloxone Tablets:

- Do not change your dose. Take Pentazocine and Naloxone Tablets exactly as prescribed by
- your healthcare provider. Use the lowest dose possible for the shortest time needed. Take your prescribed dose every 3 or 4 hours at the same time every day. Do not take more than your prescribed dose. If you miss a dose, take your next dose at your usual time.
- Call your healthcare provider if the dose you are taking does not control your pain If you have been taking Pentazocine and Naloxone Tablets regularly, do not stop taking
- Pentazocine and Naloxone Tablets without talking to your healthcare provider. Dispose of expired, unwanted, or unused Pentazocine and Naloxone Tablets by promptly flushing down the toilet, if a drug take-back option is not readily available. Visit www.fda.gov/drugdisposal for additional information on disposal of unused medicines.

# While taking Pentazocine and Naloxone Tablets DO NOT:

- Drive or operate heavy machinery, until you know how Pentazocine and Naloxone Tablets affect you. Pentazocine and Naloxone Tablets can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with Pentazocine and Naloxone Tablets may cause you to overdose and die.

# The possible side effects of Pentazocine and Naloxone Tablets:

constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are

## Get emergency medical help if you have:

trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face. tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

hese are not all the possible side effects of Pentazocine and Naloxone Tablets. Call your doctor for medical advice about side effects. You may report side effects to FDA at -800-FDA-1088. For more information go to dailymed.nlm.nih.gov.

For more information call Lupin Pharmaceuticals, Inc. at 1-866-403-7592

## This Medication Guide has been approved by the U.S. Food and Drug Administration.

08/22/2019 02:11 PM / NP Item# NOVE-NP 41212 / page 2 of 2


Manufactured by: Novel Laboratories, Inc. Somerset, NJ 08873

Manufactured for: Lupin Pharmaceuticals, Inc. Baltimore MD 21202

PI6800000211 Rev. 08/2019







## Opioid antagonists should not be administered in the absence of clinically significant respiratory or circulatory depression secondary to pentazocine overdose.

Initiate treatment with pentazocine hydrochloride and naloxone hydrochloride tablets, USP in a dosing range of 1 tablet every three to four hours. This may be increased to 2 tablets when needed. Total daily dosage


If the level of pain increases after dosage stabilization, attempt to identify the source of increased pain before

Sale Reduction or Discontinuation of Pentazocine and Natioxone Tablets in patients who may be physically dependent on opioids. Rapid discontinuation of opioid analgesics in patients who are physically dependent on opioids. Rapid discontinuation of opioid analgesics in patients who are physically dependent on opioids has resulted in serious withdrawal symptoms, uncontrolled pain, and suicide. Rapid discontinuation has also been associated with attempts to find other sources of opioid analgesics, which may be confused with drug-seeking for abuse. Patients may also attempt to treat their pain or withdrawal symptoms with illicit opioids, such as heroin, and other substances.

When a decision has been made to decrease the dose or discontinue therapy in an opioid-dependent natient

There are no standard opioid tapering schedules that are suitable for all patients. Good clinical practice dictates a patient-specific plan to taper the dose of the opioid gradually. For patients on Pentazocine and Naloxone Tablets who are physically opioid-dependent, initiate the taper by a small enough increment (e.g., no greater than 10% to 25% of the total daily dose) to avoid withdrawal symptoms, and proceed with dose-lowering at an interval of every 2 to 4 weeks. Patients who have been taking opioids for briefer periods

Dispense in a tight, light-resistant container as defined in the USP.



Store Pentazocine and Naloxone Tablets securely and dispose of properly [See PRECAUTIONS/Information

Manufactured by: **Novel Laboratories, Inc.** Somerset, NJ 08873 Manufactured for: **Lupin Pharmaceuticals, Inc.** Baltimore, MD 21202 PI6800000211

